

Intraspecific variability in paradoxid trilobites from the Purujosa trilobite assemblage (middle Cambrian, northeast Spain)

Jorge Esteve

Acta Palaeontologica Polonica 59 (1), 2014: 215-240 doi: <http://dx.doi.org/10.4202/app.2012.0006>

Eccaparadoxides? pradoanus and *Eccaparadoxides mediterraneus* are both widespread trilobite species described from the middle Cambrian of the Mediterranean region. Analysis based on a pooled sample of 500 specimens demonstrates that many of the characters that have been used to define these species show continuous variations, some of which are related to ontogeny. In addition, morphometric analyses of metric characters show that the two species cannot be distinguished on the basis of these characters either. Many of the characters studied herein are widely used in definitions and descriptions of other paradoxid species, which suggests that the taxonomic classification of other paradoxids may be oversplit.

Key words: Arthropoda, Trilobita, Paradoxididae, morphological plasticity, taphonomy, Cambrian, Murero Formation, Spain.

Jorge Esteve [jorgeves@unizar.es], Área de Paleontología, Departamento de Ciencias de la Tierra, Universidad de Zaragoza, E-50009, Zaragoza, Spain.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.