

Scanning electron microscopy of sessile graptolites

Adam Urbanek, Grażyna Mierzejewska and Piotr Mierzejewski
Acta Palaeontologica Polonica 25 (2), 1980: 197-212

The surfaces of rhabdosomes in dendroid and tuboid graptolites and in *Mastigograptus* were investigated with SEM. In general the surfaces are remarkably smooth and essentially featureless, even around the apertures. The only exception are subparallel rollers recognized on one specimen of *Dictyonema* sp., which pass at places into accumulations producing a reticulated network. The evidence available indicates a foreign nature these structures which cannot be considered units of secretion of graptolite cortex. The opinion is advanced that the recent data are suggestive of collagen fibrillogenesis occurring immediately adjacent to the surface coat of epithelial cells. The lack of bandages in sessile graptolites and the lack of foreign agglutinated material on their rhabdosomes. the intimate control of fibrillogenesis by epithelial cells in extracellular secretion of collagen are all presented as evidence in favour of the membrane model of graptolite periderm secretion.

Key words: Graptolites, scanning electron microscopy, rhabdosome surface, bandages, inclusions, collagen fibrillogenesis, models of secretion.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.