

The origin of ammonoid locomotion

Christian Klug and Dieter Korn

Acta Palaeontologica Polonica 49 (2), 2004: 235-242

Evolution of the coiled ammonoid conch from the uncoiled bactritid conch was probably coupled with changes in manoeuvrability and swimming velocity. The gradual transformation of uncoiled to coiled ammonoid conchs has essential functional consequences. The radical change in conch geometry during phylogeny but also in ontogeny of early ammonoids implies a shift of the aperture from an original roughly downward, via a downward oblique and an upward oblique to an upward orientation, presuming a neutrally buoyant condition of the ammonoid animal. Similar trends were reconstructed for the three main ammonoid lineages in the Middle Devonian, the agoniaticid, the anarcestid, and the tornoceratid lineages. This allowed an increase in manoeuvrability and in the maximum horizontal swimming speed.

Key words: Bactritida, Ammonoidea, ontogeny, phylogeny, locomotion, coiling, Devonian.

Christian Klug [chkflug@pim.unizh.ch] Paläontologisches Institut und Museum der Universität Zürich, Karl Schmid–Str. 4, 8006 Zürich, Switzerland; Dieter Korn [dieter.korn@museum.hu-berlin.de] Humboldt–Universität zu Berlin, Museum für Naturkunde, Institut für Paläontologie, D–10115 Berlin, Germany.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.