

First tillodont from India: Additional evidence for an early Eocene faunal connection between Europe and India?

Kenneth D. Rose, Rajendra S. Rana, Ashok Sahni, Kishor Kumar, Lachham Singh, and Thierry Smith
Acta Palaeontologica Polonica 54 (2), 2009: 351-355 doi:<http://dx.doi.org/10.4202/app.2008.0067>

Vastan Lignite Mine in southeastern Gujarat, India, produces the oldest known Cenozoic land-mammals and the only early Eocene continental vertebrate fauna known from India (e.g., Bajpai et al. 2005; Rana et al. 2005, 2008; Rose et al. 2006, 2008, and in press; Smith et al. 2007; Rage et al. 2008). The fauna comes from the Cambay Shale Formation and has been dated as middle Ypresian (~52 Ma, early Cuisian) based on a common nummulitid foraminiferan from about 15 m above the vertebrate-producing layer (Sahni et al. 2006; Rana et al. 2008). However, a recent study of dinoflagellate cysts from the section suggests that the deposits may be as old as 54–55 Ma (Garg et al. 2008). Although some elements of the fauna, such as anthracobunids and lagomorphs, have Asian affinities, a surprising number of taxa among the snakes, bats, insectivores, primates, rodents, and artiodactyls appear to be most closely related to early Eocene European or North American taxa. This may simply reflect the poor state of knowledge of contemporary south Asian vertebrate faunas; alternatively, it might be evidence of previously unsuspected early Eocene faunal exchange between Europe and southwest Asia. We report here two teeth of a tillodont from Vastan Mine, which constitute the first record of the mammalian order Tillodontia known from India. Despite the much greater generic diversity of tillodonts in Asia than elsewhere, the Vastan tillodont shows clear affinities with Euramerican esthonychines.

Kenneth D. Rose [kdrose@jhmi.edu], Center for Functional Anatomy and Evolution, The Johns Hopkins University School of Medicine, 1830 E Monument St., Baltimore, Maryland 21205 USA; Rajendra S. Rana [Rajendra.Rana1@gmail.com], Department of Geology, H.N.B. Garhwal University, Srinagar 246175, Uttarakhand, India; Ashok Sahni [ashok.sahni@gmail.com], Department of Geology, Panjab University, Sector 14, Chandigarh 160014, India; Kishor Kumar [kumark@wihg.res.in], Wadia Institute of Himalayan Geology, 33 General Mahadeo Singh Road, Dehradun 248001, Uttarakhand, India; Lachham Singh [rawat.lachham@yahoo.com], Department of Geology, H.N.B. Garhwal University, Srinagar 246175, Uttarakhand, India; Thierry Smith [Thierry.Smith@naturalsciences.be], Department of Paleontology, Royal Belgian Institute of Natural Sciences, B-1000 Brussels, Belgium.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(187.3 kB\)](#)