Alleged cnidarian *Sphenothallus* in the Late Ordovician of Baltica, its mineral composition and microstructure

Olev Vinn and Kalle Kirsimäe

Acta Palaeontologica Polonica 60 (4), 2015: 1001-1008 doi:<http://dx.doi.org/10.4202/app.00049.2013>

Sphenothallus is a problematic fossil with possible cnidarian affinities. Two species of *Sphenothallus*, *S. aff. longissimus* and *S. kukersianus*, occur in the normal marine sediments of the Late Ordovician of Estonia. *S. longissimus* is more common than *S. kukersianus* and has a range from early Sandbian to middle Katian. *Sphenothallus* had a wide paleo-biogeographic distribution in the Late Ordovician. The tubes of *Sphenothallus* are composed of lamellae with a homogeneous microstructure. The homogeneous microstructure could represent a diagenetic fabric, based on the similarity to diagenetic structures in *Torellella* (Cnidaria?, Hyolithelminthes). Tubes of *Sphenothallus* have an apatitic composition, but one tube contains lamellae of diagenetic calcite within the apatitic structure. *Sphenothallus* presumably had originally biomineralized apatitic tubes. Different lattice parameters of the apatite indicate that biomineralization systems of phosphatic cnidarians *Sphenothallus* and *Conularia* sp. may have been different.

Key words: Cnidaria?, *Sphenothallus*, apatite, microstructure, Ordovician, Sandbian, Katian, Estonia.

Olev Vinn [olev.vinn@ut.ee] and Kalle Kirsimäe [kalle.kirsimae@ut.ee], Department of Geology, University of Tartu, Ravila 14A, 50411 Tartu, Estonia.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.