Quail-thrush birds from the Miocene of northern Australia

Jacqueline M.T. Nguyen, Michael Archer, and Suzanne J. Hand

Acta Palaeontologica Polonica 63 (3), 2018: 493-502 doi:<https://doi.org/10.4202/app.00485.2018>

Quail-thrushes (Passeriformes: Cinclosomatidae: *Cinclosoma*) are ground-dwelling corvoid songbirds endemic to Australia and New Guinea. Until now, the only known quail-thrush fossils have been from late Quaternary cave deposits in Australia. A new species of quail-thrush, *Cinclosoma elachum* sp. nov., is described from the early to middle Miocene deposits in the Riversleigh World Heritage Area, Queensland, Australia. A second, larger quail-thrush is identified from the middle Miocene of Riversleigh. The new fossils considerably extend the geographic and temporal ranges of cinclosomatids, and indicate the presence of two species of quail-thrushes in the Miocene of northern Australia, located more than 300 km from the nearest extant member of *Cinclosoma*. These fossils provide a minimum age of ~18 million years for Cinclosomatidae. They cannot be confidently assigned to the crown group of the genus *Cinclosoma*, but can be used to calibrate the split between this genus and *Ptilorhoa* in molecular dating studies of the Corvides radiation and Passeriformes overall. This material also adds to the growing diversity of songbirds identified from the pre-Pleistocene record of Australia, and reemphasises the global significance of the Riversleigh deposits in developing understanding about the early evolutionary history of passerines.

Key words: Aves, Passeriformes, Cinclosomatidae, *Cinclosoma*, Miocene, Australia, Riversleigh.

Jacqueline M.T. Nguyen [jacqueline.nguyen@austmus.gov.au], Australian Museum Research Institute, Australian Museum, 1 William Street, Sydney, New South Wales 2010, Australia; PANGEA Research Centre, School of Biological, Earth and Environmental Sciences, UNSW Sydney, Sydney, New South Wales 2052, Australia. Michael Archer [m.archer@unsw.edu.au] and Suzanne J. Hand [s.hand@unsw.edu.au], PANGEA Research Centre, School of Biological, Earth and Environmental Sciences, UNSW Sydney, Sydney, New South Wales 2052, Australia.

distribution, and reproduction in any medium, provided the original author and source are credited.

[Full text \(252.7 kB\)](#)