Postcranial osteology of the first early-stage juvenile skeleton of *Plateosaurus trossingensis* from the Norian of Frick, Switzerland

Darius Nau, Jens N. Lallensack, Ursina Bachmann, and P. Martin Sander

Acta Palaeontologica Polonica 65 (4), 2020: 679-708 doi:<https://doi.org/10.4202/app.00757.2020>

Owing to monospecific mass-accumulation sites in Central Europe, the early-branching sauropodomorph *Plateosaurus* has one of the best fossil records among dinosaurs. Despite this, early-stage juveniles have been conspicuously absent. However, such specimens are critical in assessing the ontogenetic development of this taxon, as well as the role of heterochrony in sauropodomorph evolution. A new skeleton from the *Plateosaurus* bonebed at the Gruhalde Quarry (Klettgau Formation, Norian) of Frick, Switzerland, nicknamed “Fabian”, represents the first substantially complete juvenile referable to *Plateosaurus*. The specimen includes large portions of the cranium and vertebral column and an almost completely represented appendicular skeleton. Its juvenile ontogenetic stage is confirmed by a lack of neurocentral suture fusion in the axial skeleton. Consistent with this, the estimated total length and body mass of approximately of 2.3 m and 40 kg are considerably smaller than any previously reported specimen of the genus. Surprisingly, the postcranial morphology of the specimen is remarkably consistent with that of osteologically mature individuals, including a virtually fully developed pattern of laminae and fossae in the vertebrae. Comparisons of body proportions are complicated by varying degrees of compaction in the limb elements, but skeletal proportions mostly appear to follow isometry, with the notable exceptions of a relatively long neck, proportionately larger manus, shorter, more gracile humerus and shorter forearm in the juvenile specimen. The observed morphology suggests that adult morphology was either achieved early in ontogeny of *Plateosaurus*, or alternatively that developmental plasticity, which has previously been found to result in high variability of adult body size, could potentially also extend to morphological development.

Key words: Dinosauria, Sauropodomorpha, morphometrics, morphology, small body size, early ontogeny, Triassic, Swiss Plateau.

Darius Nau [dariusnau@uni-bonn.de], Section Paleontology, Institute of Geoscience, University of Bonn, Nussallee 8, 53115 Bonn, Germany. Jens N. Lallensack [info@dinospuren.de], Section Paleontology, Institute of Geoscience, University of Bonn, Nussallee 8, 53115 Bonn, Germany; School of Natural Sciences and Psychology, Liverpool John Moores University, James Parsons Building, Bryon

Street, Liverpool L3 3AF, UK. Ursina Bachmann [urbach@gmx.ch],
Sauriermuseum Frick, Schulstrasse 22, 5070 Frick, Switzerland. P. Martin
Sander [martin.sander@uni-bonn.de], Section Paleontology, Institute of Geoscience, University of Bonn,
Nussallee 8, 53115 Bonn, Germany; Dinosaur Institute, Natural History Museum of Los Angeles County,
900 Exposition Boulevard, Los Angeles, CA 90007, USA.

This is an open-access article distributed under the terms of the Creative Commons
Attribution License (for details please see creativecommons.org), which permits unrestricted use,
distribution, and reproduction in any medium, provided the original author and source are credited.

[Full text \(3,024.1 kB\)](#) |

[Supplementary file \(299.9 kB\)](#)