Femoral histology and growth patterns of the ceratopsian dinosaur *Psittacosaurus sibiricus* from the Early Cretaceous of Western Siberia

Pavel P. Skutschas, Semyon S. Morozov, Alexander O. Averianov, Sergey V. Leshchinskiy, Stepan V. Ivantsov, Alexey V. Fayngerts, Olga A. Feofanova, Olga N. Vladimirova and Dmitry A. Slobodin
Acta Palaeontologica Polonica 66 (2), 2021: 437-447 doi:<https://doi.org/10.4202/app.00819.2020>

The early ceratopsian dinosaur *Psittacosaurus sibiricus* from the Early Cretaceous of Western Siberia, Russia, is one of the most advanced and largest (up to 2.5 m) members of the genus. Here we present a description of ontogenetic changes in the long-bone histology of this species. Analysis of a growth series of femora demonstrates significant histological maturation during ontogeny, expressed by the progressive appearance of signs of bone remodeling (erosion bays, secondary bone formation), decreasing of vascularity, changing of the orientation of vascular canals from reticular to longitudinal, and appearance of parallel-fibred bone in the outer part cortex. These ontogenetic changes in the longbone histology of *P. sibiricus* are generally similar to those of another relatively advanced species, *P. lujiatunensis* from China. The basal *P. mongoliensis* from Mongolia shows less mature long-bone histology during late ontogeny (e.g., the late appearance of signs of remodeling and the predominance of reticular vascularization at later stages). We suggest that the earlier achievement of histological maturity is an evolutionary trend of the genus *Psittacosaurus*.

Key words: Dinosauria, Ceratopsia, *Psittacosaurus*, bone histology, Cretaceous, Siberia, Russia.

Pavel P. Skutschas [skutchas@mail.ru; p.skutschas@spbu.ru] and Semyon S. Morozov [semyon-morozow7@yandex.ru], Vertebrate Zoology Department, Saint Petersburg State University, Universitetskaya nab. 7/9, 199034 Saint Petersburg, Russia. Alexander O. Averianov [dzharakuduk@mail.ru], Zoological Institute of the Russian Academy of Sciences, Universitetskaya nab. 1, 199034 Saint Petersburg, Russia. Stepan V. Ivantsov [stepan_ivantsov@mail.ru], Laboratory of Micropaleontology, Tomsk State University, Prospekt Lenina 36, 634050, Tomsk, Russia. Sergey V. Leshchinskiy [sl@ggf.tsu.ru] and Alexey V. Fayngerts [alex@ggf.tsu.ru], Laboratory of Mesozoic and Cenozoic Continental Ecosystems, Tomsk State University, Prospekt Lenina 36, 634050, Tomsk, Russia. Olga A. Feofanova [feofanova42@yandex.ru], Olga N. Vladimirova [ovlad-5@mail.ru], and Dmitry A. Slobodin [d.a.slobodin@mail.ru], Kuzbass State Museum of Local Lore, Prospekt Sovetskiy 51, 650000, Kemerovo, Russia.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[Full text \(3,901.8 kB\)](#)