

Siliceous sponges from the Upper Cretaceous of Poland Part 1. Tetraxonia

Helena Hurcewicz

Acta Palaeontologica Polonica 11 (1), 1966: 15-129

Sixty two species of sponges, belonging to the Tetraxonia, of this number, 19 new ones: *Brochodora latiramea* n.sp., *Pachypoterion biedai* n.sp., *Homalodora skrainivensis* n.sp., *H. polonica* n.sp., *H. brachiramosa* n.sp., *Heterostinia phytoniformis* n.sp., *Heloraphinia chordata* n.sp., *Phymatella irregularis* n.sp., *Thecosiphonia gracilis* n.sp., *Kozlowskispongia bulbosa* n.gen., n.sp., *Phyllodermia magna* n.sp., *Ph. pulchra* n.sp., *Eustrobilus extraneus* n.sp., *Ragadinia foraminifera* n.sp., *Plinthosella elegans* n.sp., *Acrochordonia regularis* n.sp. and *A. bifurcata* n.sp. are described and one new genus, *Kozlowskispongia* n.gen., is erected. The species investigated belong to Tetractinellida (2 species), Tetracladina (32 species), Megacladina (16 species) and Dicranocladina (7 species). In most species, the morphology of the body, the structure of the skeleton and the morphology of the megascleres have been studied. In many of them, the microscleres have been found.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.