

## Lower Permian bryozoans from southern and central Spitsbergen, Svalbard

Hans Arne Nakrem, Błażej Błażejowski, and Andrzej Gaździcki

*Acta Palaeontologica Polonica* 54 (4), 2009: 677-698 doi: <http://dx.doi.org/10.4202/app.2008.0078>

Bryozoans from the Lower Permian Treskelodden and Wordiekammen formations of southern and central Spitsbergen

respectively, Svalbard, have been studied. Twenty species are identified, including one new genus, *Toulapora* gen. nov., with *Toulapora svalbardense* as type species and one new species, *Ascopora birkenmajeri* sp. nov. The taxonomic composition is typical Lower Permian, with species in common with Timan–Pechora and the Urals (Russia) and Ellesmere Island (the Canadian Arctic). Growth habits reflect a moderately to deeper shelf environment.

**Key words:** Bryozoa, Permian, Spitsbergen, Arctic.

Hans Arne Nakrem [[h.a.nakrem@nhm.uio.no](mailto:h.a.nakrem@nhm.uio.no)], University of Oslo, Natural History Museum, PO Box 1172 Blindern, 0318 Oslo, Norway; Błażej

Błażejowski [[bblazej@twarda.pan.pl](mailto:bblazej@twarda.pan.pl)] and Andrzej Gaździcki [[gazdzick@twarda.pan.pl](mailto:gazdzick@twarda.pan.pl)] Instytut Paleobiologii PAN, Twarda 51/55, 00–818 Warszawa, Poland

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.