Lower Permian bryozoans from southern and central Spitsbergen, Svalbard

Hans Arne Nakrem, Błażej Błażejowski, and Andrzej Gaździcki

Acta Palaeontologica Polonica 54 (4), 2009: 677-698 doi: <http://dx.doi.org/10.4202/app.2008.0078>

Bryozoans from the Lower Permian Treskelodden and Wordiekammen formations of southern and central Spitsbergen

respectively, Svalbard, have been studied. Twenty species are identified, including one new genus, *Toulapora* gen. nov., with *Toulapora svalbardense* as type species and one new species, *Ascopora birkenmajeri* sp. nov. The taxonomic composition is typical Lower Permian, with species in common with Timan–Pechora and the Urals (Russia) and Ellesmere Island (the Canadian Arctic). Growth habits reflect a moderately to deeper shelf environment.

Key words: Bryozoa, Permian, Spitsbergen, Arctic.

Hans Arne Nakrem [h.a.nakrem@nhm.uio.no], University of Oslo, Natural

History Museum, PO Box 1172 Blindern, 0318 Oslo, Norway; Błażej

Błażejowski [bblazej@twarda.pan.pl] and Andrzej Gaździcki [gazdzick@twarda.pan.pl] Instytut

Paleobiologii PAN, Twarda 51/55, 00–818 Warszawa, Poland

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.