Shallow-water occurrence of *Wiwaxia* in the middle Cambrian of the Barrandian area, Czech Republic

Oldřich Fatka, Petr Kraft, and Michal Szabad

Acta Palaeontologica Polonica 56 (4), 2011: 871-875 doi: <http://dx.doi.org/10.4202/app.2009.0052>

Isolated sclerites of the genus *Wiwaxia* are reported from shale interlayers in lower levels of middle Cambrian (unnamed 3rd Series of Cambrian) Buchava Formation in the Skryje–Týřovice Basin in the Czech Republic interpreted as shallow–water sediments. Geographic distribution of *Wiwaxia* indicates latitudinal control as all occurrences are obviously restricted to tropical belt.

Oldřich Fatka [fatka@natur.cuni.cz] and Petr Kraft [kraft@natur.cuni.cz], Charles University, Institute of Geology and Palaeontology, Albertov 6, 128 43 Praha 2, Czech Republic; Michal Szabad [geosz.pb@seznam.cz], Obránců míru 75, 261 02 Příbram VII, Czech Republic.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.