Morphology and ontogeny of the Cambrian edrioasteroid echinoderm *Cambraster cannati* from western Gondwana

Samuel Zamora, Colin D. Sumrall, and Daniel Vizcaïno

Acta Palaeontologica Polonica 58 (3), 2012: 545-559 doi: <http://dx.doi.org/10.4202/app.2011.0152>

A review of the Cambrian edrioasteroid echinoderm *Cambraster cannati* is made based on new collections from the Iberian Chains (NE Spain) and Montagne Noire (France). New morphological data include a completely articulated oral area and details of ambulacra. Specimens ranging from 4 to 26 mm in diameter provide detailed information concerning the full ontogeny. Important changes through ontogeny mainly affect the marginal ring and the plating pattern of the aboral surface. Comparison with other species of *Cambraster* indicates that the aboral surface of *Cambraster tastudorum* from Australia shows strong resemblance to juvenile specimens of *C. cannati*. *Cambraster cannati* was attached directly to the substrate and inhabited relatively high energy, offshore environments from the west margin of Gondwana. Abnormalities in the skeleton are described for the first time in a Cambrian edrioasteroid.

Key words: Echinodermata, Edrioasteroidea, palaeobiology, Spain, France.

Samuel Zamora [samuel@unizar.es], Department of Palaeontology, The Natural History Museum, Cromwell Road, London SW7 5BD, UK; Colin D. Sumrall [csumrall@utk.edu], Department of Earth and Planetary Sciences, University of Tennessee, Knoxville, TN 37996, USA; Daniel Vizcaïno [daniel.vizcaino@wanadoo.fr] 7 rue J.-B. Chardin, Maquens, 11090 Carcassonne, France.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[Full text \(1,812.3 kB\)](#)