Taxonomy and palaeobiogeography of the Cenozoic Euro-Mediterranean rissoid gastropod *Galeodinopsis* and its relationship with close genera

Vittorio Garilli and Daniela Parrinello

Acta Palaeontologica Polonica 59 (2), 2014: 379-406 doi: <http://dx.doi.org/10.4202/app.2012.0044>

The investigation of the Late Paleogene to Late Neogene species of rissoid gastropod *Galeodinopsis* in the Euro-Mediterranean area has supported the hypothesis that this genus is an intermediate form between two well-known rissoids, *Alvinia* and *Manzonina*. We recognized four species of *Galeodinopsis*: *G. biangulata*, *G. germanica* sp. nov., *G. semperi* (new name for *Rissoa duboisii*), and *G. tiberiana*. The oldest (very Late Eocene/Oligocene) representatives of *Galeodinopsis*, *G. biangulata* and *G. semperi*, share similar shell shape and microsculpture with *Alvinia*. This suggests that *Galeodinopsis* originated from some Eocene species related to *Alvinia*. The new species represents the first occurrence within *Galeodinopsis* of a combination of characters very close to those of *Manzonina*, above all the typical pitted microsculpture. We hypothesize that *Manzonina* evolved from *Galeodinopsis* rather than from the genera *Alvinia* or *Alvinia*, as previously supposed. *Galeodinopsis* originated during the very Late Eocene/Oligocene in the North Sea Basin. Afterwards it underwent a strong southward shift to the mid-high east Atlantic and the Mediterranean area, during the Mio-Pliocene, and to the Recent tropical eastern Atlantic coasts, where the type species *G. tiberiana* still lives. The shift likely was due to a combination of climate cooling and palaeogeographical changes. The distribution of *G. biangulata* suggests that connections between the North Sea Basin and the Atlantic domain opened through the Channel area at least during the Early Oligocene, earlier than indicated previously. The distribution of *Manzonina moulinsi* supports the idea of a southern connection to the Atlantic Aquitaine Basin via the Rhine Graben during the Late Oligocene. From a palaeoecological point of view, *Galeodinopsis* includes warm species with planktotrophic larval development that are typical of the shelf environment in fully marine conditions.

Key words: Gastropoda, Rissoidae, taxonomy, palaeogeography, biogeography, Cenozoic, North Sea, Atlantic connection, Europe.

Vittorio Garilli [vittoriogarilli@apema.eu], APEMA Research and Educational Service, Via Alla Falconara 34, 90136 Palermo, Italy; Daniela Parrinello [daniela.parrinello@unipa.it], Facoltà di Scienze della Formazione, University of Palermo, Via Archirafi 18, 90100 Palermo, Italy.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[Full text \(2.008.1 kB\)](#) |

[Supplementary file \(103.5 kB\)](#)