

Autothecal morphs and dormancy in the camaroid graptolite *Xenotheka*

Piotr Mierzejewski

Acta Palaeontologica Polonica 48 (1), 2003: 93-98

The camaroid graptolite *Xenotheka klinostoma* Eisenack, 1937 is described from the lower Llanvirn limestones of Gilbergabrottet, northern Öland, Sweden. Two distinct autothecal morphs are recognized: (1) normal morph (described for the first time), i.e. an autotheca with an unsculptured outer surface, devoid of both an outer lining and autothecal occlusion, and inhabited by an active zooid; and (2) sealed morph, i.e. an autotheca coated and occluded, provided with a sculptured outer lining made of a unique verrucose fabric, and inhabited by an inactive or dormant zooid. In addition, the existence of a hypothetical (3) unsealed morph or re-opened autotheca, devoid of an autothecal occlusion but provided with an outer lining, and inhabited by a reactivated zooid, is predicted. The sealed morphs may represent an adaptation which allowed their inhabitants to survive adverse conditions. The outer lining of *Xenotheka* is compared with a peculiar outer membrane found in the modern hemichordate *Rhabdopleura*, from the intertidal zone of Fiji, and with camaroid extracamaral tissue.

Key words: Graptolithina, Camaroidea, microfossils, ultrastructure, dormancy, Ordovician, Sweden.

Piotr Mierzejewski [mierzejewski@graptolite.net], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland; present address: ul. Filtrowa 83 m. 49, PL-02-032 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.