

## A new helmeted frog of the genus *Thaumastosaurus* from the Eocene of England

J. Alan Holman and David L. Harrison

*Acta Palaeontologica Polonica* 48 (1), 2003: 157-160

Helmeted frogs with co-ossified skulls (*Thaumastosaurus*) were previously known in Europe only in the Eocene on the basis of *Thaumastosaurus bottii* from southwestern France and *Thaumastosaurus wardi* from coastal southern England. We describe *Thaumastosaurus sulcatus*, that differs from these species in having a different dermal sculpture pattern and tooth morphology. We suggest that the auburn colored tooth crown tips in *T. sulcatus*, as in some other vertebrates, reflect extra hardness for protection against tooth wear. Autochthonous *Thaumastosaurus* species in southern England support the concept that southern England and south-western France were distinct faunal provinces in the Late Eocene.

J. Alan Holman [[holman@msu.edu](mailto:holman@msu.edu)], Michigan State University Museum, East Lansing, Michigan, 48824–1045, USA; David L. Harrison [[hzm@btinternet.com](mailto:hzm@btinternet.com)], Harrison Institute, St. Botolphs Road, Sevenoaks, Kent, TN3–3AQ, England

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.