

Growth and life habits of the Triassic cynodont *Trirachodon*, inferred from bone histology

Jennifer Botha and Anusuya Chinsamy

Acta Palaeontologica Polonica 49 (4), 2004: 619-627

Growth pattern and lifestyle habits of the Triassic non-mammalian cynodont *Trirachodon* are deduced from bone histology and cross-sectional geometry. Several skeletal elements of *Trirachodon* were examined in order to document histological changes during ontogeny, as well as histovariability in the skeleton. The bone histology of all the elements consists of a moderately vascularized, periodically interrupted, fibro-lamellar bone tissue. This suggests that the overall growth of *Trirachodon* was probably rapid during the favourable season, but decreased or ceased during the unfavourable season. As the environment is thought to have been semi-arid with seasonal rainfall, it is possible that *Trirachodon* was sensitive to such environmental fluctuations. Some inter-elemental histovariability was noted where the number and prominence of growth rings varied. Limb bone cross-sectional geometry revealed a relatively thick bone wall and supports earlier proposals that *Trirachodon* was fossorial.

Key words: Cynodonts, *Trirachodon*, lifestyles, bone histology, growth patterns.

Jennifer Botha [jbotha@iziko.org.za], Natural History Collections Division, South African Museum, Iziko Museums of Cape Town, P.O. Box 61, 8000, South Africa (corresponding author); Anusuya Chinsamy [achinsam@botzoo.uct.ac.za], Zoology Department, University of Cape Town, Rondebosch, 7701, South Africa.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

