

Precursory siphuncular membranes in the body chamber of *Phyllopachyceras* and comparisons with other ammonoids

Kazushige Tanabe, Cyprian Kulicki, and Neil H. Landman


Acta Palaeontologica Polonica 50 (1), 2005: 9-18

Organic membranes preserved in the rear part of the body chamber of the Late Cretaceous phylloceratid ammonite *Phyllopachyceras ezoense* were examined with scanning electron microscopy (SEM) on the basis of well-preserved specimens from Hokkaido, Japan. SEM observations revealed that the membranes are continuous with the siphuncular tube wall in the phragmocone and consist of two layers, both of which are made of a dark, primarily conchiolin material; namely, a thinner inner homogeneous layer and a thicker outer layer with gently inclined pillar-like units. Hence, they are interpreted as the precursory siphuncular membranes. The precursory siphuncular membranes are not associated with any other organic components such as the siphuncular sheets reported in some Paleozoic and Mesozoic ammonoids. Unlike the tube-like condition in the phragmocone, the precursory siphuncular membranes in the body chamber of the specimens examined do not form a tube shape; on the ventral side the membranes are truncated and directly contact the outer shell wall. These observations suggest that the inner and outer layers of the precursory siphuncular membranes in the body chamber were respectively formed by the siphuncular epithelium from the inner side and by the invaginated septal epithelium from the outer side. It is also postulated that at the initial stage of septal formation, the rear part of the body moved slowly forward, developing a circum-siphonal invagination of the septal epithelium. Because similar conchiolin membranes are occasionally preserved in the body chambers of other phylloceratids, the above morphogenetic process applies to all members of the Phylloceratina. The tube-shaped structure in the rear part of the body chamber of desmoceratid *Damesites* consists only of nacreous layer. We interpret it as a pathologically overgrown prochoanitic septal neck.

Key words: Ammonoidea, *Phyllopachyceras*, morphogenesis, conchiolin membranes, siphuncle, septal epithelium, chamber formation.

Kazushige Tanabe [tanabe@eps.s.u-tokyo.ac.jp], Department of Earth and Planetary Science, University of Tokyo, Tokyo 113-0033, Japan; Cyprian Kulicki [kulicki@twarda.pan.pl], Instytut Paleobiologii, Polska Akademia Nauk, ul. Twarda 51/55, PL-00-818 Warszawa, Poland; Neil H. Landman [landman@amnh.org], Division of Paleontology (Invertebrates), American Museum of Natural History, Central Park West at 79th Street, New York, New York 10024-5192, USA.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,932.1 kB\)](#)