

Pyritized tube-feet in a protasterid ophiuroid from the Upper Ordovician of Kentucky, U.S.A.

Alexander Glass Acta Palaeontologica Polonica 51 (1), 2006: 171-184

A single specimen of the protasterid ophiuroid Protasterina flexuosa from the Kope Formation (Cincinnatian, Upper Ordovician) of Kentucky exhibits three-dimensionally pyritized tube feet. This represents the first report of soft-tissue preservation in an echinoderm from the type-Cincinnatian series. The tube feet are solid and lack all internal structure. They consist of aggregated masses of small euhedral to subhedral pyrite crystals suggesting that pyritization, although decay-induced and mediated, did not necessarily replicate soft-tissues but might instead have formed inside the void-spaces left behind during the decay process. The discovery of pyritized soft-tissue as delicate as ophiuroid tube feet suggests that similar forms of soft-tissue preservation might be found in other taxa in the Kope Formation. Perhaps much more importantly, this unexpected occurrence demonstrates the incompleteness of our knowledge of permissible conditions for the preservation of soft-tissues and it thereby indicates promise for discovery of other such occurrences in diverse organisms in unexpected settings. Systematics of Paleozoic ophiuroids remains problematic in spite of many years of study by capable paleontologists. The incomplete but well-preserved specimens treated here include the types of Protasterina flexuosa and Protasterina fimbriata as well as previously undescribed specimens. Together they permit a revised diagnosis and detailed description of the genus Protasterina. Protasterina fimbriata is the type species of the genus but is a subjective junior synonym of *Protaster flexuosus* (= *Protasterina flexuosa*). The genus is clearly differentiated from the only other known protasterid ophiuroid from the Cincinnatian series, Taeniaster spinosus, and from all other protasterid genera.

Key words: Ophiuroidea, Protasteridae, soft-tissue preservation, pyritization, Cincinnatian, Edenian, Ordovician, Kentucky.

Alexander Glass <u>glassa@geology.cwu.edu</u>, Department of Geological Sciences, Central Washington University, 400 East University Avenue, Ellensburg, WA 98926-7418, United States of America.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

