

## Early Cretaceous multituberculate mammals from the Kuwajima Formation (Tetori Group), central Japan

Nao Kusuhashi

*Acta Palaeontologica Polonica* 53 (3), 2008: 379-390 doi:<http://dx.doi.org/10.4202/app.2008.0302>

*Hakusanobaatar matsuoi* gen. et sp. nov. and *Tedoribaatar reini* gen. et sp. nov. are multituberculate mammals recovered from the Lower Cretaceous (Barremian to lower Aptian) Kuwajima Formation of the Tetori Group in the Shiramine district, Hakusan City, Ishikawa Prefecture, central Japan. *Hakusanobaatar matsuoi* is an eobaatarid multituberculate characterized by a P4 with cusp formula 3:5, and a P5 with cusp formula 2:6:2. One of the specimens of *H. matsuoi* has the best preserved upper premolar series among known eobaatarid specimens. Based on the dentition of *H. matsuoi*, it is highly probable that the cimolodontan P4 is homologous with the “plagiaulacidan” P5. *Tedoribaatar reini* is also tentatively attributed to Eobaataridae, and shows a single-rooted p3 and loss of at least the permanent p2. On the basis of these apomorphic features, *T. reini* is considered to be the “plagiaulacidan” multituberculate that is most closely related to cimolodontans.

**Key words:** Mammalia, Multituberculata, Eobaataridae, Hakusanobaatar, Tedoribaatar, Early Cretaceous, Kuwajima Formation, Tetori Group, Japan

Nao Kusuhashi [[nao\\_kusuhashi@ivpp.ac.cn](mailto:nao_kusuhashi@ivpp.ac.cn)], Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044, People's Republic of China, and Department of Geology and Mineralogy, Graduate School of Science, Kyoto University, Kyoto 606–8502, Japan.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

