

The titanosaur sauropods from the late Campanian-early Maastrichtian Allen Formation of Salitral Moreno, Río Negro, Argentina

Rodolfo A. Garcia and Leonardo Salgado *Acta Palaeontologica Polonica* 58 (2), 2013: 269-284 doi: http://dx.doi.org/10.4202/app.2011.0055

The dinosaur record of the Salitral Moreno locality (Río Negro Province, Argentina) is characterized by a high diversity of herbivore taxa, among them hadrosaurs, ankylosaurs, and titanosaur sauropods, but carnivores are rare, consisting of only a few fragmentary bones of small forms. Titanosaurs are represented by *Rocasaurus muniozi* and *Aeolosaurus* sp., and at least four other taxa, represented by fragmentary material. The elements preserved include a cervical, dorsal and caudal vertebrae, chevron, humerii, ulnae, radii, metacarpal, femora, tibiae, metatarsal, ischia, pubis, and ilium. The Allen Formation is thought to be correlated with the Marília Formation in Brazil, and their faunas have certain elements in common such as aeolosaurines, but saltasaurines and hadrosaurs, are known exclusively from the Allen Formation. These absences, and particularly that of the saltasaurines, may be because those sauropods originated late in the Cretaceous, probably in southern South America (Northern Patagonia?), and they did not have time to disperse to northern South America.

Key words: Saurischia, Sauropodomorpha, Sauropoda, Titanosauria, Titanosaur, Salitral Moreno, Allen Formation, Maastrichtian, Patagonia.

Rodolfo A. García [rodosnow@yahoo.com.ar] and Leonardo Salgado [salgadoleito@yahoo.com.ar], Conicet-Inibioma, Museo de Geología y Paleontología Universidad Nacional del Comahue, Buenos Aires 1400, Neuquén (8300), Argentina.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

