

Fauna and stratigraphy of the Uppermost Triassic and the Toarcian and Aalenian deposits in Sassenfjorden, Spitsbergen

Andrzej Wierzbowski, Cyprian Kulicki and Halina Pugaczewska

Acta Palaeontologica Polonica 26 (3-4), 1981: 195-237

Five sections of the De Geerdalen Formation (mostly the *Wilhelmeya* Member) and the Janusfjellet Formation (the Brentskardhaugen Bed) between Janusfjellet and Botneheia in Sassenfjorden (Spitsbergen) were studied stratigraphically and palaeontologically. The fauna of the *Wilhelmsya* Member indicates an uppermost Triassic age. The fauna from phosphatic nodules in the condensed Brentskardhaugen Bed is of Middle Toarcian to Early Aalenian age. Pelecypods assigned to 18 genera are described as well as ammonites representing five genera. *Harpoceras kopiki* Wierzbowski et Kulicki sp.n. is established.

Key words: Upper Triassic, Toarcian, Aalenian, biostratigraphy, ammonites, pelecypods, Spitsbergen.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.