

## Multituberculate mammals from the Cretaceous of Uzbekistan

Zofia Kielan- Jaworowska and Lev A. Nesson

*Acta Palaeontologica Polonica* 37 (1), 1992: 1-17

The first western Asian multituberculates found in the Bissekty Formation (Coniacian) of Uzbekistan are described on the basis of a lower premolar (p4), a fragment of a lower incisor, an edentulous dentary, a proximal part of the humerus and a proximal part of the femur. *Uzbekbaatar kizylkumensis* gen. et sp. n. is defined as having a low and arcuate p4, possibly without a posterobuccal cusp; it presumably had two lower premolars, as inferred from the presence of a triangular concavity at the upper part of the anterior wall of p4, and p3 less reduced in relation to p4 than in non-specialized Taeniolabidoidea and Ptilodontoidea. *Uzbekbaatar* is placed in the Cimolodonta without indicating family and infraorder. It might have originated from the Plagiaulacinae or Eobaatarinae.

**Key words:** Multituberculata, Mammalia, Cretaceous, Coniacian, Uzbekistan

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,009.8 kB\)](#)