

## Palaeobiological and palaeoenvironmental significance of the Pliocene trace fossil *Dactyloidites peniculus*

Alfred Uchman and Peter Pervesler

*Acta Palaeontologica Polonica* 52 (4), 2007: 799-808

The radial trace fossil *Dactyloidites peniculus* occurs in a deep tier in totally bioturbated shoreface sediments of Pliocene age in the Stirone Valley, N Italy, together with *Thalassinoides* isp. and *Ophiomorpha nodosa*. Long, narrow shafts running from centre of the radiating structure and abundant faecal pellets in the radial structure were discovered. The trace maker of *D. peniculus*, probably a polychaete, deposited the pellets deeply in the sediment, probably for reinforcement of the tubes and a gardening of microbes for feeding. This trace fossil exclusively occurs within a narrow horizon at the top of a shallowing-up section interpreted as a high-stand system tract, below a discontinuity surface capped by finer sediments. *D. peniculus* was formed in soft sandy sediments under stable conditions related to the latest phases of the high-stand system tract. Therefore, it is a candidate for indication of similar environmental situations having a soft sandy, but stable sea floor.

**Key words:** Trace fossils, *Dactyloidites*, gardening, sequence stratigraphy, marine flooding, foredeep, Neogene, Pliocene.

Alfred Uchman [fred@geos.ing.uj.edu.pl](mailto:fred@geos.ing.uj.edu.pl) Institute of Geological Sciences, Jagiellonian University, ul. Oleandry 2a, PL-30-063 Kraków, Poland; Peter Pervesler [peter.pervesler@univie.ac.at](mailto:peter.pervesler@univie.ac.at) Department of Palaeontology, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

