



http://app.pan.pl/SOM/app68-Simon_Salgado_SOM.pdf

SUPPLEMENTARY ONLINE MATERIAL FOR

A new gigantic titanosaur (Dinosauria, Sauropoda) from the early Late Cretaceous of Patagonia (Neuquén Province, Argentina)

María Edith Simón and Leonardo Salgado

Published in *Acta Palaeontologica Polonica* 2023 68 (4): 719-735.

<https://doi.org/10.4202/app.01086.2023>

Supplementary Online Material

SOM 1. List of characters based on Pérez Moreno *et al.* (2022) and character scoring for *Bustingorrytitan shiva* gen. et sp. nov.

SOM 2. Autapomorphic characters of *Bustingorrytitan shiva* gen. et sp. nov recovered in the analysis

Bibliography

Table 1. Measurements of the caudal vertebrae of *Bustingorrytitan shiva* gen. and sp. nov.

Table 2. Measurements of the humeri of *Bustingorrytitan shiva* gen. and sp. nov.

Table 3. Measurements of the ulnae of *Bustingorrytitan shiva* gen. and sp. nov.

Table 4. Measurements of the metacarpals of *Bustingorrytitan shiva* gen. and sp. nov.

Table 5. Measurements of the femora of *Bustingorrytitan shiva* gen. and sp. nov.

Table 6. Measurements of the tibiae of *Bustingorrytitan shiva* gen. and sp. nov.

Table 7. Measurements of the metatarsals of *Bustingorrytitan shiva* gen. and sp. nov.

Table 8. Measurements of the ungual phalanges of *Bustingorrytitan shiva* gen. and sp. nov.

Matrix data available at http://app.pan.pl/SOM/app68-Simon_Salgado_SOM/matrix.tnt

SOM 1. List of characters based on Pérez Moreno *et al.* (2022) and character scoring for *Bustingorrytitan shiva* gen. et sp. nov.

Skull

1. Posterolateral processes of premaxilla and lateral processes of maxilla, shape: without midline contact (0); with midline contact forming marked narial depression, subnarial foramen not visible laterally (1). (Wilson, 2002:character 1). ***Bustingorrytitan*=?**
2. Premaxillary anterior margin shape: without step (0); with marked step but short step (1); with marked and long step (2) (modified from Wilson, 2002:character 2). ***Bustingorrytitan*=?**
3. Premaxila, ascending process shape in lateral view: convex (0); concave, with a large dorsal projection (1); sub-rectilinear and directed posterodorsally (2). (Whitlock, 2011:character 3) ***Bustingorrytitan*=?**
4. Premaxilla, external surface: without anteroventrally orientated vascular grooves originating from an opening in the maxillary contact (0); vascular grooves present (1). (Whitlock, 2011:character 2) ***Bustingorrytitan*=?**
5. Premaxilla-maxilla suture, shape: planar (0); twisted along its length, giving the contact a sinuous appearance in lateral view (1). (D'Emic, 2012:character 2) ***Bustingorrytitan*=?**
6. Premaxilla, small finger-like, vertically oriented premaxillary process near anteromedial corner of external naris: (0) absent; (1) present. (D'Emic, 2012:character 3) ***Bustingorrytitan*=?**
7. Maxillary border of external naris, length: short, making up much less than onefourth narial perimeter (0); long, making up more than one third narial perimeter (1). (Wilson, 2002:character 3). ***Bustingorrytitan*=?**
8. Maxilla, foramen anterior to the preantorbital fenestra: absent (0); present (1). (Zaher *et al.*, 2011:character 244). ***Bustingorrytitan*=?**
9. Preanteorbital fenestra: absent (0); present, being wide and laterally opened (1). (Modified from Wilson, 2002:character 4). ***Bustingorrytitan*=?**
10. Subnarial foramen and anterior maxillary foramen, position: well distanced from one another (0); separated by narrow bony isthmus (1). (Wilson, 2002:character 5) ***Bustingorrytitan*=?**
11. Antorbital fenestra: much shorter than orbital maximum diameter, less than 85% of orbit (0); subequal to orbital maximum diameter, greater than 85% orbit (1). (Modified from Wilson, 2002:character 6 following to Whitlock, 2011:character 13) ***Bustingorrytitan*=?**

12. Antorbital fenestra, shape of dorsal margin: straight or convex (0); concave (1). (Whitlock, 2011:character 14). ***Bustingorrytitan*=?**
13. Antorbital fossa: present (0); absent (1). (Wilson, 2002:character 7) ***Bustingorrytitan*=?**
14. External nares position: terminal (0); retracted to level of orbit (1); retracted to a position between orbits (2). (Wilson, 2002:character 8) ***Bustingorrytitan*=?**
15. External nares, maximum diameter: shorter (0); or longer than orbital maximum diameter (1). (Wilson, 2002:character 9) ***Bustingorrytitan*=?**
16. Orbital ventral margin, anteroposterior length: broad, with subcircular orbital margin (0); reduced, with acute orbital margin (1). (Wilson, 2002:character 10) ***Bustingorrytitan*=?**
17. Lacrimal, anterior process: present (0); absent (1). (Wilson, 2002:character 11) ***Bustingorrytitan*=?**
18. Lacrimal, anteriorly projecting vertical plate of bone: absent (0); present (1). (D'Emic, 2012:character 4) ***Bustingorrytitan*=?**
19. Jugal contribution to the ventral border of the skull: present and long (0); absent or very reduced (1). (Carballido *et al.*, 2012:character 16) ***Bustingorrytitan*=?**
20. Quadratojugal-Maxilla contact: absent or small (0); broad (1). (Whitlock, 2011:character 10) ***Bustingorrytitan*=?**
21. Jugal-ectopterygoid contact: present (0); absent (1). (Wilson, 2002:character 12) ***Bustingorrytitan*=?**
22. Jugal, contribution to antorbital fenestra: absent (0); present, but very reduced (1); present and large, bordering approximately one-third its perimeter (2). (Modified from Wilson, 2002:character 13) ***Bustingorrytitan*=?**
23. Quadratojugal, position of anterior terminus: posterior to orbit (0); between the orbit (1); anteriorly surpassing the orbit (2). (modified by Canudo *et al.*, 2018 in the base of Whitlock, 2011:character 30) ***Bustingorrytitan*=?**
24. Quadratojugal, anterior process length: short, anterior process shorter than dorsal process (0); long, anterior process more than twice as long as dorsal process (1). (Wilson, 2002:character 32) ***Bustingorrytitan*=?**
25. Quadratojugal, angle between anterior and dorsal processes: less than or equal to 90°, so that the quadrate shaft is directed dorsally (0); greater than 90°, approaching 130°, so that the quadrate shaft slants posterodorsally (1). (Whitlock, 2011:character 31) ***Bustingorrytitan*=?**

26. Ventral edge of anterior surface of the quadratojugal: straight, not expanded ventrally (0); slightly expanded ventrally, forming a small bulge, which height is less than twice the ramus height (1); well expanded ventrally, forming a notorious bulge, which height is twice or more the minimum height of the ramus (2).
 (Modified from Upchurch *et al.*, 2004:character 26) **Bustingorrytitan=?**
27. Squamosal contribution to the supratemporal fenestra: present, the squamosal is well visible in dorsal view (0); reduced or absent (1). (Curry Rogers, 2005:character 37) **Bustingorrytitan=?**
28. Squamosal-quadratojugal contact: present (0); absent (1). (Wilson, 2002:character 31) **Bustingorrytitan=?**
29. Squamosal, posteroventral margin: smooth (0); "with prominent, ventrally directed ""prong"" (1). (Whitlock, 2011:character 37) **Bustingorrytitan=?**
30. Prefrontal posterior process size: small, not projecting far posterior of frontal-nasal suture (0); elongate, approaching parietal (1). (Wilson, 2002:character 14) **Bustingorrytitan=?**
31. Prefrontal, posterior process shape: flat (0); hooked (1). (Wilson, 2002:character 15) **Bustingorrytitan=?**
32. Prefrontal, anterior process: absent (0); present (1). (Curry Rogers, 2005:character 30) **Bustingorrytitan=?**
33. Prefrontal-Frontal contact width: large, equal or longer than the anteroposterior length of the prefrontal (0); narrow, less than half the anteroposterior length of the prefrontal (1). (Zaher *et al.*, 2011:character 239)
Bustingorrytitan=?
34. Postorbital, ventral process shape: transversely narrow (0); broader transversely than anteroposteriorly (1).
 (Wilson, 2002:character 16) **Bustingorrytitan=?**
35. Postorbital, posterior process: present (0); absent (1). (Wilson, 2002:character 17) **Bustingorrytitan=?**
36. Postorbital, posterior margin articulating with the squamosal: with tapering posterior process (0); with a deep posterior process (1). (Zaher *et al.*, 2011:character 245) **Bustingorrytitan=?**
37. Frontal contribution to supratemporal fossa: present (0); absent (1). (Wilson, 2002:character 18)
Bustingorrytitan=?
38. Frontals, midline contact (symphysis): sutured (0); or fused in adult individuals (1). (Wilson, 2002:character 19) **Bustingorrytitan=?**
39. Frontal, anteroposterior length: approximately twice (0); or less than minimum transverse breadth (1).
 (Wilson, 2002:character 20) **Bustingorrytitan=?**

40. Frontal-nasal suture, shape: flat or slightly bowed anteriorly (0); V-shaped, pointing posteriorly (1). (Whitlock, 2011:character 21) **Bustingorrytitan=?**
41. Frontals, dorsal surface: without paired grooves facing anterodorsally (0); grooves present, extend on to nasal (1). (Whitlock, 2011:character 22) **Bustingorrytitan=?**
42. Frontal, contribution to dorsal margin of orbit: contribution to dorsal margin of orbit: less than 1.5 times the contribution of prefrontal (0); at least 1.5 times the contribution of prefrontal (1). (Whitlock, 2011:character 23) **Bustingorrytitan=?**
43. Parietal occipital process, dorsoventral height: short, less than the diameter of the foramen magnum (0); deep, nearly twice the diameter of the foramen magnum (1). (Wilson, 2002:character 21) **Bustingorrytitan=?**
44. Parietal, contribution to post-temporal fenestra: present (0); absent (1). (Wilson, 2002:character 22)
Bustingorrytitan=?
45. Parietal, distance separating supratemporal fenestrae: less than the long axis of supratemporal fenestra, 0.8 or less (0); almost the same than the long axis of supratemporal fenestra 0.8-1.2 (1); much larger than the long axis of supratemporal fenestra more than 1.2 (2). (Modified from Wilson, 2002:character 24)
Bustingorrytitan=?
46. Postparietal foramen: absent (0); present (1). (Wilson, 2002:character 23) **Bustingorrytitan=?**
47. Paroccipital process distal terminus: straight, slightly expanded surface (0); rounded, tongue-like process (1). (Whitlock, 2011:character 42) **Bustingorrytitan=?**
48. Supratemporal fenestra: present (0); absent (1). (Wilson, 2002:character 25) **Bustingorrytitan=?**
49. Supratemporal fenestra, long axis orientation: anteroposterior (0); transverse (1). (Wilson, 2002:character 26)
Bustingorrytitan=?
50. Supratemporal fenestra, maximum diameter: much longer than (0); or subequal to that of foramen magnum (1). (Wilson, 2002:character 27) **Bustingorrytitan=?**
51. Supratemporal region, anteroposterior length: temporal bar longer (0); or shorter anteroposteriorly than transversely (1). (Wilson, 2002:character 28) **Bustingorrytitan=?**
52. Supratemporal fossa, lateral exposure: not visible laterally, obscured by temporal bar (0); visible laterally, temporal bar shifted ventrally (1). (Wilson, 2002:character 29) **Bustingorrytitan=?**

53. Supraoccipital, sagital nuchal crest: broad, weakly developed (0); narrow, sharp and distinct (1). (Whitlock, 2011:character 45) ***Bustingorrytitan***=?
54. Laterotemporal fenestra, anterior extension: posterior to orbit (0); ventral to orbit (1). (Wilson, 2002:character 30) ***Bustingorrytitan***=?
55. Quadratojugal fossa: absent (0); present (1). (Wilson, 2002:character 33) ***Bustingorrytitan***=?
56. Quadratojugal fossa, depth: shallow (0); deeply invaginated (1). (Wilson, 2002:character 34) ***Bustingorrytitan***=?
57. Quadratojugal fossa, orientation: posterior (0); posterolateral (1). (Wilson, 2002:character 35) ***Bustingorrytitan***=?
58. Quadratojugal, articular surface shape: quadrangular in ventral view, oriented transversely or roughly triangular in shape (0); thin, crescent-shaped surface with anteriorly directed medial process (1). (Whitlock, 2011:character 32) ***Bustingorrytitan***=?
59. Quadratojugal, articular surface shape: quadrangular in ventral view, oriented transversely or roughly triangular in shape (0); thin, crescent-shaped surface with anteriorly directed medial process (1). (Whitlock, 2011:character 32) ***Bustingorrytitan***=?
60. Palatobasal contact, shape: pterygoid with small facet (0); dorsomedially orientated hook (1); or rocker-like surface for basipterygoid articulation (2). (Wilson, 2002:character 36) ***Bustingorrytitan***=?
61. Pterygoid, transverse flange (i.e. ectopterygoid process) position: posterior of orbit (0); between orbit and antorbital fenestra (1); anterior to antorbital fenestra (2). (Wilson, 2002:character 37) ***Bustingorrytitan***=?
62. Pterygoid, quadrate flange size: large, palatobasal and quadrate articulations well separated (0); small, palatobasal and quadrate articulations approach (1). (Wilson, 2002:character 38) ***Bustingorrytitan***=?
63. Pterygoid, palatine ramus shape: straight, at level of dorsal margin of quadrate ramus (0); stepped, raised above level of quadrate ramus (1). (Wilson, 2002:character 39) ***Bustingorrytitan***=?
64. Pterygoid, sutural contact with ectopterygoid: broad, along the medial or lateral surface (0); narrow, restricted to the anterior tip of the ectopterygoid (1). (Zaher *et al.*, 2011:character 240) ***Bustingorrytitan***=?
65. Palatine, lateral ramus shape: plate-shaped (long maxillary contact) (0); rod-shaped (narrow maxillary contact) (1). (Wilson, 2002:character 40) ***Bustingorrytitan***=?
66. Epopterygoid: present (0); absent (1). (Wilson, 2002:character 41) ***Bustingorrytitan***=?
67. Vomer, anterior articulation: maxilla (0); premaxilla (1). (Wilson, 2002:character 42) ***Bustingorrytitan***=?

68. Supraoccipital, height: twice subequal to (0); or less than height of foramen magnum (1). (Wilson, 2002:character 43) ***Bustingorrytitan*=?**
69. Paroccipital process, ventral non-articular process: absent (0); present (1). (Wilson, 2002:character 44) ***Bustingorrytitan*=?**
70. Crista prootica, size: rudimentary (0); expanded laterally into dorsolateral process (1). (Wilson, 2002:character 45) ***Bustingorrytitan*=?**
71. Basipterygoid processes, length: short, approximately twice (0); or elongate, at least four times basal diameter (1). (Wilson, 2002:character 46) ***Bustingorrytitan*=?**
72. Basipterygoid processes, angle of divergence: approximately 45° (0); less than 30° (1). (Wilson, 2002:character 47) ***Bustingorrytitan*=?**
73. Basal tubera, anteroposterior depth: approximately half dorsoventral height (0); sheet-like, 20% dorsoventral height (1). (Wilson, 2002:character 48) ***Bustingorrytitan*=?**
74. Basal tubera, breadth: much broader than (0); or narrower than occipital condyle (1). (Wilson, 2002:character 49) ***Bustingorrytitan*=?**
75. Basal tubera: distinct from basipterygoid (0); reduced to slight swelling on ventral surface of basipterygoid (1). (Whitlock, 2011:character 53) ***Bustingorrytitan*=?**
76. Basal tubera, shape of posterior face: convex (0); slightly concave (1). (Whitlock, 2011:character 54) ***Bustingorrytitan*=?**
77. Basioccipital depression between foramen magnum and basal tubera: absent (0); present (1). (Wilson, 2002:character 50) ***Bustingorrytitan*=?**
78. Basisphenoid/basipterygoid recess: present (0); absent (1). (Wilson, 2002:character 51) ***Bustingorrytitan*=?**
79. Basisphenoid/quadrat contact: absent (0); present (1). (Wilson, 2002) ***Bustingorrytitan*=?**
80. Basisphenoid, sagital ridge between basipterygoid processes: absent (0); present (1). (Zaher *et al.*, 2011:character 242) ***Bustingorrytitan*=?**
81. Basipterygoid processes, orientation: perpendicular to (0); or angled approximately 45° to skull roof (1). (Wilson, 2002:character 53) ***Bustingorrytitan*=?**

82. Basipterygoid, area between the basipterygoid processes and parasphenoid rostrum: is a mildly concave subtriangular region (0); forms a deep slot-like cavity that passes posteriorly between the bases of the basipterygoid processes (1). (Mannion *et al.*, 2013:character 48) **Bustingorrytitan=?**
83. Occipital region of skull, shape: anteroposteriorly deep, paroccipital processes oriented posterolaterally (0); flat, paroccipital processes oriented transversely (1). (Wilson, 2002:character 54) **Bustingorrytitan=?**
84. Occipital condyle, lateral surface of the basioccipital: flat or slightly convex (0); strongly concave (1). (Remes *et al.*, 2009:character 50) **Bustingorrytitan=?**
85. Dentary, depth of anterior end of ramus: slightly less than that of dentary at midlength (0); 150% minimum depth (1). (Wilson, 2002:character 55) **Bustingorrytitan=0**
86. Dentary, anteroventral margin shape: gently rounded (0); sharply projecting triangular process (1). (Wilson, 2002:character 56) **Bustingorrytitan=0**
87. Dentary symphysis, orientation: angled 15° or more anteriorly to (0); or perpendicular to axis of jaw ramus (1). (Wilson, 2002:character 57) **Bustingorrytitan=1**
88. Dentary, cross-sectional shape of symphysis: oblong or rectangular (0); subtriangular, tapering sharply towards ventral extreme (1); subcircular (2). (Whitlock, 2011:character 60) **Bustingorrytitan=0**
89. Dentary, tuberosity on labial surface near symphysis: absent (0); present (1). (Whitlock, 2011:character 57) **Bustingorrytitan=0**
90. Dentary, posteroventral process shape: single (0); divided (1). (D'Emic, 2012:character 10)
Bustingorrytitan=?
91. Mandible, coronoid eminence: strongly expressed, clearly rising above plane of dentigerous portion (0); absent (1). (Whitlock, 2011:character 62) **Bustingorrytitan=?**
92. External mandibular fenestra: present (0); absent (1). (Wilson, 2002:character 58) **Bustingorrytitan=?**
93. Surangular depth: less than twice (0); or more than two and one-half times maximum depth of the angular (1). (Wilson, 2002:character 59) **Bustingorrytitan=?**
94. Surangular ridge separating adductor and articular fossae: absent (0); present (1). (Wilson, 2002:character 60)
Bustingorrytitan=?
95. Adductor fossa, medial wall depth: shallow (0); deep, prearticular expanded dorsoventrally (1). (Wilson, 2002:character 61) **Bustingorrytitan=?**

96. Splenial posterior process, position: overlapping angular (0); separating anterior portions of prearticular and angular (1). (Wilson, 2002:character 62) ***Bustingorrytitan=?***
97. Splenial posterodorsal process: present, approaching margin of adductor chamber (0); absent (1). (Wilson, 2002:character 63) ***Bustingorrytitan=?***
98. Coronoid, size: extending to dorsal margin of jaw (0); reduced, not extending dorsal to splenial (1); absent (2). (Wilson, 2002:character 64) ***Bustingorrytitan=?***
99. Tooth rows, shape of anterior portions: narrowly arched, anterior portion of tooth rows V-shaped (0); broadly arched, anterior portion of tooth rows U-shaped (1); rectangular, tooth-bearing portion of jaw perpendicular to jaw rami (2). (Wilson, 2002:character 65). ***Bustingorrytitan=?***
100. Tooth rows, length: extending to orbit (0); restricted anterior to orbit (1); restricted anterior to antorbital fenestra (2); restricted anterior to subnarial foramen (3). (Modified from Wilson, 2002:character 66)
Bustingorrytitan=?
101. Maxillary teeth shape: straight along axis (0); twisted axially through an arc of 30- 45°: absent (0); present (1). (D'Emic, 2012:character 15) ***Bustingorrytitan=?***
102. Dentary teeth, number: greater than 20 (0); 10-17 (1); 9 or fewer (2). (Modified from Wilson, 2002:character 73) ***Bustingorrytitan=?***
103. Replacement teeth per alveolus, number: two or fewer (0); more than four (1). (Wilson, 2002:character 74)
Bustingorrytitan=0
104. Lateral plate: absent (0); present (1). (Upchurch *et al.*, 2004:character 9) ***Bustingorrytitan=?***
105. Teeth, orientation: perpendicular (0); or oriented anteriorly relative to jaw margin (1). (Wilson, 2002:character 75) ***Bustingorrytitan=0***
106. Tooth crowns, orientation: aligned along jaw axis, crowns do not overlap (0); aligned slightly anterolingually, tooth crowns overlap (1). (Wilson, 2002:character 69) ***Bustingorrytitan=?***
107. Tooth crowns, shape: narrow crowns (0); broad crowns (1). ***Bustingorrytitan=0***
108. Tooth crowns, cross-sectional shape at mid-crown: elliptical (0); D-shaped (1); subcylindrical (2); cylindrical (3). (Wilson, 2002:character 70) ***Bustingorrytitan=1***
109. SI values for tooth crowns: less than 3.0 (0); 3.0-4.0 (1); 4.0-5.0 (2); more than 5.0 (3). (Upchurch *et al.*, 2004:chs. 67-69) ***Bustingorrytitan=?***

110. Crown-to-crown occlusion: absent (0); present (1). (Wilson, 2002:character 67) **Bustingorrytitan=0**

111. V-shaped wear facets: present (0); absent (1). (Modified from Wilson, 2002:character 68)

Bustingorrytitan=?

112. Development of the marginal wear facets: well developed (0); slightly developed as marginal facets (1)

Bustingorrytitan=?

113. One high angle wear facet and a second low angle wear facet: absent (0); present (1) ***Bustingorrytitan=?***

114. Single planar wear facet in labial or lingual surface of the teeth: absent (0); present (1) ***Bustingorrytitan=?***

115. Marginal tooth denticles: present (0); absent on posterior edge (1); absent on both anterior and posterior edges (2). (Wilson, 2002:character 72) ***Bustingorrytitan=2***

116. Enamel surface texture: smooth (0); wrinkled (1). (Wilson, 2002:character 71) ***Bustingorrytitan=0***

117. Thickness of enamel asymmetric labiolingually: absent (0); present (1). (Whitlock, 2011:character 74)

Bustingorrytitan=0

118. Teeth, longitudinal grooves on lingual aspect: absent (0); present (1). (Wilson, 2002:character 76)

Bustingorrytitan=0

Cervical vertebrae

119. Cervical vertebrae, number: 10 or fewer (0); 12 (1); 13-14 (2); 15 (3); 16 or more (4). (Modified from Wilson, 2002:character 80 and Upchurch *et al.*, 2004:chs. 96-100) ***Bustingorrytitan=?***

120. Atlas, intercentrum occipital facet shape: rectangular in lateral view, length of dorsal aspect subequal to that of ventral aspect (0); expanded anteroventrally in lateral view, anteroposterior length of dorsal aspect shorter than that of ventral aspect (1). (Wilson, 2002:character 79) ***Bustingorrytitan=?***

121. Axis, centrum shape: over two and a half times as long as tall (0); less than twice as long as tall (1). (D'Emic, 2012:character 20) ***Bustingorrytitan=?***

122. Cervical vertebrae, parapophyses, shape and orientation: short and weakly developed, projected laterally or slightly ventrally (0); middle development, ventrally such that the cervical ribs are displaced ventrally around half the height of the centrum (1); well developed, broad and ventrally projected such that cervical ribs are displaced ventrally more than the height of the centrum (2). (Modified from D'Emic, 2012:character 29)

Bustingorrytitan=0

123. Cervical centra, articulations: amphicoelous (0); opisthocoelous (1). (Salgado *et al.*, 1997:character 1 ; Wilson, 2002:character 82; Upchurch, 1998:character 81 and Upchurch *et al.*, 2004:character 103)

Bustingorrytitan=1

124. Cervical centra, ventral surface: is flat or slightly convex transversely (0); transversely concave (1). (Upchurch, 1998:character 84 and Upchurch *et al.*, 2004:character 107) **Bustingorrytitan=1**
125. Cervical centra, midline keels on ventral surface: prominent and plate-like (0); reduced to low ridges or absent (1). (Upchurch, 1998:character 83 and Upchurch *et al.*, 2004:character 106) **Bustingorrytitan=1**
126. Cervical centra, pleurocoels: absent (0); present with well defined anterior, dorsal, and ventral edges, but not the posterior one (1); present, with well defined edges (2); absent, but with deep lateral fossa which bears small pneumatopores that communicate to the interior pneumatic cavities (3) **Bustingorrytitan=0**
127. Cervical centra, pleurocoels: singles without division (0); with a well defined anterior excavation and a posterior smooth fossa (1); divided by a bone septum, resulting in an anterior and a posterior lateral excavation (2); divided in three or more lateral excavations, resulting in a complex morphology (3); with a well defined anterior excavation and a posterior smooth fossa (Modified from Salgado *et al.*, 1997; Wilson, 2002; Harris, 2006) **Bustingorrytitan=NA**
128. Cervical vertebrae, well-developed epiphyses: absent (0); present (1). **Bustingorrytitan=?**
129. Cervical vertebrae, epiphyses shape: stout, pillar-like expansions above postzygapophyses (0); posteriorly projecting prongs (1). (D'Emic, 2012:character 24) **Bustingorrytitan=?**
130. Prezygapophyses, anterior process suited ventrolaterally to the articular surface: absent (0); present (1). (Remes *et al.*, 2009:character 79) **Bustingorrytitan=?**
131. Cervical vertebrae with an accessory lamina, which runs from the PODL (or slightly anteriorly) up to the SPOL: absent (0); present (1). (Modified from D'Emic, 2012:character 25) **Bustingorrytitan=?**
132. Cervical vertebrae, height divided width (measured in its posterior articular surface): higher than 1.1 (0), around 1 (1); between 0.9 and 0.7 (2); smaller than 0.7 (3). (Modified from Wilson, 2002:character 84; Upchurch, 1998:character 85 and Upchurch *et al.*, 2004:character 108) **Bustingorrytitan=?**
133. Cervical centra, small notch in the dorsal margin of the posterior articular surface: absent (0); present (1). (Carballido *et al.*, 2012) **Bustingorrytitan=?**

134. Cervical vertebrae, neural arch lamination: well developed, with well-marked laminae and fossae (0); rudimentary, with diapophyseal laminae absents or very slightly marked (1). (Wilson, 2002:ch, 81)

Bustingorrytitan=?

135. Cervical vertebrae with an accessory lamina, which runs from the postzygodiaiapophyseal lamina (PODL) up to the spinoprezygapophyseal lamina (SPRL): absent (0); present (1). (Modified from Sereno *et al.*, 2007:characters 50 and 51; Whitlock, 2011:characters 78 and 96) **Bustingorrytitan=0**

136. Cervical centra, internal pneumaticity: absent (0); present with singles and wide cavities (1); present, with several small and complex internal cavities (2). (Modified from Carballido *et al.*, 2011) **Bustingorrytitan=?**

137. Anterior cervical vertebrae, prespinal lamina: absent (0); present (1). (Carballido *et al.*, 2012)

Bustingorrytitan=0

138. Anterior cervical vertebrae, neural spine shape: single (0); bifid (1). (Wilson, 2002:character 72; Upchurch *et al.*, 2004:character 118) **Bustingorrytitan=?**

139. Middle and posterior cervical vertebrae, prespinal lamina: absent (0); present (1). (Carballido *et al.*, 2012)

Bustingorrytitan=?

140. Middle cervical vertebrae, lateral fossae on the prezygapophysis process: absent (0); present (1). (Harris, 2006) **Bustingorrytitan=?**

141. Middle, cervical vertebrae, height of the neural arch: less than the height of the posterior articular surface (0); higher than the height of the posterior articular surface (1). (Wilson, 2002:character 87; similar Upchurch *et al.*, 2004:111 and 112) **Bustingorrytitan=?**

142. Middle cervical centrum, anteroposterior length divided the height of the posterior articular surface: less than 4 (0); more than 4 (1). (Wilson, 2002:character 74; and Upchurch *et al.*, 2004:character 102)

Bustingorrytitan=?

143. Middle and posterior cervical vertebrae, morphology of the centroprezygapophyseal lamina: single (0); dorsally divided, resulting in a lateral and medial lamina, being the medial lamina linked with the intraprezygapophyseal lamina and not with the prezygapophysis (1); divided, resulting in the presence of a “true” divided centroprezygapophyseal lamina, which is dorsally connected to the prezygapophysis (2).

(Carballido *et al.*, 2012) **Bustingorrytitan=?**

144. Middle and posterior cervical vertebrae, morphology of the centropostzygapophyseal lamina (CPOL): single (0); divided, with the medial part contacting the intrapostzygapophyseal lamina (1) (Carballido *et al.*, 2012)

Bustingorrytitan=?

145. Middle and posterior cervical vertebrae, articular surface of zygapophyses: flat (0); transversally convex (1). (Upchurch *et al.*, 2004) **Bustingorrytitan=?**

146. Middle and posterior cervical vertebrae, prominent triangular flange on the posterior edge of the diapophyseal process (in the PCDL): absent (0); present (1). (Remes *et al.*, 2009:character 78)

Bustingorrytitan=?

147. Middle cervical vertebrae, prezygapophyses position: do not extend beyond the anterior margin of the centrum (0); extends beyond the anterior margin of the centrum (1). (Salgado *et al.*, 1997:character 37)

Bustingorrytitan=?

148. Middle and posterior cervical vertebrae, parapophysis shape: subcircular (0); anteroposteriorly elongate (1). (D'Emic, 2012:character 28) **Bustingorrytitan=?**

149. Posterior cervical vertebrae, lateral profile of the neural spine: displays steeply sloping cranial and caudal faces (0); displays steeply sloping cranial face and noticeably less steep caudal margin (1). (Upchurch *et al.*, 2004:character 119) **Bustingorrytitan=?**

150. Posterior cervical vertebrae, neural spine shape: not expanded distally (0); expanded but not as much as the width of the centrum (1); laterally expanded, being equal or wider than the vertebral centrum (1). (Modified from González Riga *et al.*, 2009) **Bustingorrytitan=?**

151. Posterior cervical vertebrae, lateral expansion: SPRLs does not contact the lateral margins of the neural spine (0); SPRLs are contacting the lateral margins of the neural spine (1). (Modified from González Riga and Ortiz, 2014:character 26-27) **Bustingorrytitan=?**

152. Posterior cervical and anterior dorsal vertebrae, neural spine shape: single (0); bifid (1). (Wilson, 2002:character 90, Upchurch *et al.*, 2004:character 118) **Bustingorrytitan=?**

153. Posterior cervical vertebrae, proportions – ratio total height / centrum length: less than 1.5 (0); more than 1.5 (1). (González Riga *et al.*, 2009:character 32) **Bustingorrytitan=?**

154. Posterior cervical and anterior dorsal bifid neural spines, median tubercle: absent (0); present (1)

Bustingorrytitan=?

Dorsal vertebrae

155. Number of dorsal vertebrae: 14 or more (0); 13 (1); 12 (2); 10 (3). (Modified from Wilson, 2002:character 91; Upchurch *et al.*, 2004:character 122- 125) **Bustingorrytitan=?**
156. Dorsal centra, pleurocoels: absent (0); present (1). (Wilson, 2002:character 78; Upchurch *et al.*, 2004:128)
Bustingorrytitan=1
157. Dorsal vertebrae, transverse processes: are directed laterally or slightly upwards (0); are directed strongly dorsolaterally (1). (Upchurch *et al.*, 2004:character 138) **Bustingorrytitan=?**
158. Dorsal vertebrae, distal end of the transverse process: curves smoothly into the dorsal surface of the process (0); is set off from the dorsal surface, the latter having a distinct dorsally facing flattened area (1). (Upchurch *et al.*, 2004:character 140) **Bustingorrytitan=?**
159. Anterior dorsal vertebrae, non-bifid neural spine in anterior or posterior view: posses subparallel lateral margins (0); posses lateral margins which slightly diverge dorsally (1); posses lateral margins which strongly diverge dorsally (2); posses lateral margins dorsomedially inclined, that approach each other (3). (Modified 52 from Wilson, 2002:character 107; Upchurch *et al.*, 2004:character 155) **Bustingorrytitan=?**
160. Middle to posterior dorsal vertebrae, non-bifid neural spine in anterior or posterior view: posses subparallel lateral margins (0); posses lateral margins which slightly diverge dorsally (1); posses lateral margins which strongly diverge dorsally (2). (Modified from Wilson, 2002:character 107; Upchurch *et al.*, 2004:character 155) **Bustingorrytitan=0**
161. Dorsal centra, pneumatic structures: absent, dorsal centra with solid internal structure (0); present, dorsal centra with simple and big air-spaces (camerate) (1); present, dorsal centra with small and complex air-spaces (polycamerate) (2); present, dorsal centra with small and complex air spaces (semicamellate/camellate) (3). (Modified from Carballido *et al.*, 2011) **Bustingorrytitan=0**
162. Anterior and middle dorsal neural spines, spinoprezygapophyseal lamina (SPRL): absent (0); present (1). (Modified from Upchurch *et al.*, 2007:character 131) **Bustingorrytitan=?**
163. Posterior dorsal neural spines, spinoprezygapophyseal lamina (SPRL): absent (0); present (1). (Modified from Upchurch *et al.*, 2007:character 132) **Bustingorrytitan=?**

164. Dorsal vertebrae, single not bifid neural spines, single prespinal lamina (PRSL): absent (0); present (1).
 (Modified from Salgado *et al.*, 1997:character14) **Bustingorrytitan=1**
165. Dorsal vertebrae, single not bifid neural spines, single prespinal lamina (PRSL): rough and wide, present in the dorsalmost part of the neural spine (0); rough and wide, extended through almost all the neural spine (1); smooth and narrow (2). (Carballido *et al.*, 2012) **Bustingorrytitan=1**
166. Dorsal vertebrae with single neural spines, middle single fossa projected through the midline of the neural spine: present (0); absent (1). (Carballido *et al.*, 2012) **Bustingorrytitan=1**
167. Dorsal vertebrae with single neural spines, middle single fossa, projected through the midline of the neural spine: relatively wide median simple fossa (0); a thin median simple fossa (1); extremely reduced median simple fossa (2). (Carballido *et al.*, 2012) **Bustingorrytitan=NA**
168. Anterior dorsal centra, articular face shape: amphicoelous (0); opisthocoelous (1). (Wilson, 2002:character 94; Upchurch *et al.*, 2004:character 104) **Bustingorrytitan=?**
169. Anterior and middle dorsal centra, pleurocoels: have rounded caudal margins (0); have tapering, acute caudal margins (1). (Salgado *et al.*, 1997; Upchurch, 1998:character 06; Upchurch *et al.*, 2004:character 127)
Bustingorrytitan=?
170. Middle dorsal neural arches in lateral view, anterior edge of the neural spine: project anteriorly to the diapophysis (0); converge with the diapophysis (1); project posteriorly to the diapophysis (2). (Carballido *et al.*, 2012) **Bustingorrytitan=?**
171. Anterior and middle dorsal vertebrae, zygapophyseal articulation angle: horizontal or slightly posteroventrally oriented (0); posteroventrally oriented (around 30°) (1); strongly posteroventrally oriented (more than 40°) (2). (Carballido *et al.*, 2012) **Bustingorrytitan=?**
172. Anterior dorsal vertebrae, neural spine orientation: vertical, or slightly inclined (less than 20°) (0); posterodorsally, more than 20° (1); anteriorly directed (2) **Bustingorrytitan=?**
173. Anterior dorsal vertebrae neural spine, triangular aliform processes: absent (0); present but do not project far laterally (not as far as caudal zygapophyses) (1); present and project far laterally (as far as caudal zygapophyses) (2). (Modified from Wilson, 2002:character 102 and Upchurch *et al.*, 2004:characters 153 and 154) **Bustingorrytitan=?**

174. Anterior dorsal vertebrae, neural spine minimums width / length: 0.5 or greater (stout and short neural spine) (0); lower than 0.5 (thin and tall neural spines). (Carballido *et al.*, 2017:character 174) ***Bustingorrytitan=?***
175. Anterior dorsal vertebrae, neural spine length (from TPRL to top): less than the height of the centrum (0); slightly higher than the centrum (1); twice or more the height of the centrum (2). (Carballido *et al.*, 2017:character 175) ***Bustingorrytitan=?***
176. Anterior dorsal vertebrae, dorsal edge of the neural spine: flat (0); arrow-shaped (1); convex (2). (Carballido *et al.*, 2017:character 176) ***Bustingorrytitan=?***
177. Posterior dorsal vertebrae, dorsal edge of the neural spine: flat (0); arrow-shaped (1); convex (2). (Carballido *et al.*, 2017:character 177) ***Bustingorrytitan=?***
178. Middle to posterior dorsal centra, ventral surface: convex transversely (0); flattened (1); is slightly concave, sometimes with one or two crests (2). (Upchurch *et al.*, 2004) ***Bustingorrytitan=?***
179. Middle dorsal vertebrae, hypophene-hypantrum system: present (0); absent (1). (Modified from Salgado *et al.*, 1997:character 25; Wilson, 2002:character 106; Upchurch *et al.*, 2004:character 145) ***Bustingorrytitan=?***
180. Posterior dorsal vertebrae, hypophene-hypantrum system: present and well developed, usually with a rhomboid shape (0); present and weakly developed, mainly as a laminar articulation (1); absent or only present in posteriormost dorsal vertebrae (2). (Carballido *et al.*, 2012) ***Bustingorrytitan=0***
181. Middle and posterior dorsal vertebrae, transverse processes length: short (0); long (projecting along 1.5 the articular surface width) (1). (Carballido *et al.*, 2012) ***Bustingorrytitan=?***
182. Mid and posterior dorsal vertebrae with a single lamina (the single TPOL) supporting the hypophene or postzygapophysis from below: absent (0); present (1). (Modified from Upchurch *et al.*, 2004:character 146) ***Bustingorrytitan=1***
183. Middle and posterior dorsal vertebrae, neural canal in anterior view: entirely surrounded by the neural arch (0); enclosed in a deep fossa, enclosed laterally by pedicels (1). (Upchurch *et al.*, 2004:character 136) ***Bustingorrytitan=0***
184. Middle and posterior dorsal vertebrae, neural spine height: approximately twice the centrum length (0); for times the centrum length (1). (Upchurch *et al.*, 2004) ***Bustingorrytitan=?***

185. Middle and posterior dorsal neural spines orientation: vertical (0); slightly inclined, with an angle of around 70 degrees (1); strongly inclined, with an angle not bigger than 40 degrees (2). (Modified from Wilson, 2002:character 104) **Bustingorrytitan=?**
186. Middle and posterior dorsal vertebral, central keel: absent (0); present (1). (D'Emic, 2012:character 49)
Bustingorrytitan=?
187. Dorsal vertebrae, height of the neural arch divided the height of the centrum: less than 0.8 (0); more than 0.8 (1). (Pol *et al.*, 2011:character 132) **Bustingorrytitan=?**
188. Middle to posterior dorsal vertebrae, pleurocoel dorsal margin: rounded (0); angular (1). (Rauhut *et al.*, 2015:character 346) **Bustingorrytitan=1**
189. Middle to posterior dorsal vertebrae, pleurocoel dorsal margin: well below the dorsal margin of the centrum (0); at the level of the dorsal margin of the centrum or higher (1). (Rauhut *et al.*, 2015:character 347)
Bustingorrytitan=1
190. Middle to posterior dorsal vertebrae, small fossa anterior or anteroventral to the pleurocoel: absent (0); present (1). (Rauhut *et al.*, 2015:character 348) **Bustingorrytitan=0**
191. Middle and posterior dorsal neural arches, centropostzygapophyseal lamina (CPOL), shape: simple (0); divided (1). (Wilson, 2002:character 95) **Bustingorrytitan=1**
192. Middle and posterior dorsal neural arches, anterior centroparapophyseal lamina (ACPL): absent (0); present (1). (Wilson, 2002:character 96; Upchurch *et al.*, 2004:character 133) **Bustingorrytitan=1**
193. Middle and posterior dorsal neural arches, prezygoparapophyseal lamina (PRPL): absent (0); present (1). (Wilson, 2002:character 97) **Bustingorrytitan=1**
194. Middle and posterior dorsal neural arches, posterior centroparapophyseal lamina (PCPL): absent (0); present (1). (Wilson, 2002:character 98; Upchurch *et al.*, 2004:character 137) **Bustingorrytitan=1**
195. Middle and posterior dorsal centrum in transverse section (height: width ratio): subcircular (ratio, similar to 1 or a bit higher) (0); slightly dorsoventrally compressed (ratios between 0.8 and 1) (1); strongly compressed (ratios below 0.8) (2). (Modified from Upchurch *et al.*, 2004) **Bustingorrytitan=2**
196. Middle and posterior dorsal vertebrae neural spine, triangular aliform processes: absent (0); present but do not project far laterally (not as far as caudal zygapophyses) (1); present and project far laterally (as far as

caudal zygapophyses) (2). (Modified from Wilson, 2002:character 102 and Upchurch *et al.*, 2004:characters 153 and 154) ***Bustingorrytitan=?***

197. Middle and posterior dorsal vertebrae, spinodiapophyseal lamina (SPDL): absent (0); present (1). (Upchurch *et al.*, 2004:character 157) ***Bustingorrytitan=1***
198. Middle and posterior dorsal vertebrae, accessory spinodiapophyseal lamina (SPDL): absent (0); present (1). (Upchurch *et al.*, 2004:character 151) ***Bustingorrytitan=1***
199. Dorsal vertebrae, spinodiapophyseal webbing: lamina follows curvature of neural spine in anterior view (0); lamina "festooned" from spine, dorsal margin does not closely follow shape of neural spine and diapophysis (1). (Whitlock, 2011:character 104) ***Bustingorrytitan=?***
200. Anterior dorsal vertebrae, spinopostzygapophyseal lamina (SPOL): absent (0); present (1). (Upchurch *et al.*, 2007:character 133) ***Bustingorrytitan=?***
201. Middle and posterior dorsal neural spines, lateral spinopostzygapophyseal lamina (lSPOL): absent (0); present (1). (Wilson, 2002:character 100; Upchurch *et al.*, 2004:character 159) ***Bustingorrytitan=0***
202. Middle and posterior dorsal neural arches, spinodiapophyseal lamina (SPDL) and spinopostzygapophyseal lamina (lSPOL) contact: absent (0); present (1). (Wilson, 2002:character 101) ***Bustingorrytitan=NA***
203. Middle and posterior dorsal vertebrae, spinodiapophyseal (SPDL) and spinopostzygapophyseal lamina (lSPOL) contact: ventral, well separated from the triangular aliform process (0); dorsal, forms part of the triangular aliform process (1). (Carballido *et al.*, 2012) ***Bustingorrytitan=NA***
204. Middle and posterior dorsal vertebrae, height of neural arch below the postzygapophyses (pedicel): less than height of centrum (0); subequal to or greater than height of centrum (1). (Whitlock, 2011:character 109)
Bustingorrytitan=1
205. Posterior Dorsal vertebrae, medial spinopostzygapophyseal lamina (mSPOL): absent (0); present and forms part of the median posterior lamina (1). (Carballido *et al.*, 2012) ***Bustingorrytitan=0***
206. Posterior dorsal vertebrae, transverse processes: lie posterior, or posterodorsal, to the parapophysis (0); lie vertically above the parapophysis (1). (Upchurch *et al.*, 2004:character 139) ***Bustingorrytitan=1***
207. Posterior dorsal centra, articular face shape: amphicoelous (0); slightly opisthocoelous (1); opisthocoelous (2). (Modified from Wilson, 2002:character 105) ***Bustingorrytitan=1***

208. Posterior dorsal vertebrae, neural spine: narrower transversely than anteroposteriorly (0); broader transversely than anteroposteriorly (1). (Wilson, 2002: character 92) ***Bustingorrytitan*=?**
209. Posterior dorsal vertebra, posterior centrodiapophyseal lamina (PCDL): has an unexpanded ventral tip (0); expands and may bifurcate toward its ventral tip (1). (Salgado *et al.*, 1997) ***Bustingorrytitan*=1**

Ribs

210. Cervical ribs, distal shafts of longest cervical ribs: are elongate and form overlapping bundles (0); are short and do not project beyond the caudal end of the centrum to which they are attached (1). (Wilson, 2002:character 140) ***Bustingorrytitan*=?**
211. Cervical ribs, angle between the capitulum and tuberculum: greater than 90°, so that the rib shaft lies close to the ventral edge of the centrum (0); less than 90°, so that the rib shaft lies below the ventral margin of the centrum (1). (Wilson, 2002:character 139) ***Bustingorrytitan*=?**
212. Dorsal ribs, proximal pneumatopores: absent (0); present (1). (Wilson, 2002:character 141)
***Bustingorrytitan*=?**
213. Anterior dorsal ribs, cross-sectional shape: subcircular (0); plank-like, anteroposterior breadth more than three times mediolateral breadth (1). (Wilson, 2002) ***Bustingorrytitan*=?**

Sacrum

214. Sacral vertebrae, number: 3 or fewer (0); 4 (1); 5 (2); 6 (3). (Wilson, 2002:character 108)
***Bustingorrytitan*=?**
215. Sacrum, sacricostal yoke: absent (0); present (1). (Wilson, 2002:character 109) ***Bustingorrytitan*=?**
216. Sacral vertebrae contributing to acetabulum: numbers 1-3 (0); numbers 2-4 (1). (Wilson, 2002:character 110)
***Bustingorrytitan*=?**
217. Sacral neural spines length: approximately twice length of centrum (0); approximately four times length of centrum (1). (Wilson, 2002:character 111) ***Bustingorrytitan*=?**
218. Sacral ribs, dorsoventral length: low, not projecting beyond dorsal margin of ilium (0); high extending beyond dorsal margin of ilium (1). (Wilson, 2002:character 112) ***Bustingorrytitan*=?**

219. Pleurocoels in the lateral surfaces of sacral centra: absent (0); present (1). (Upchurch *et al.*, 2004:character 165) ***Bustingorrytitan=?***

Caudal vertebrae

220. Caudal vertebrae, number: 35 or fewer (0); 40 to 55 (1); increased to 70-80 (2). (Wilson, 2002:character 114)

Bustingorrytitan=?

221. Caudal bone texture: solid (0); spongy (camellate), with large internal cells (1). (Wilson, 2002:character 113)

Bustingorrytitan=0

222. Anterior caudals, pneumatized neural arch: absent (0); present (1). ***Bustingorrytitan=0***

223. Caudal transverse processes: persist through caudal 20 or more posteriorly (0); disappear by caudal 15 (1); disappear by caudal 10(2). (Wilson, 2002:character 115) ***Bustingorrytitan=?***

224. First caudal centrum anterior articular surface: flat (0); concave (1); convex (2). ***Bustingorrytitan=?***

225. First caudal centrum, posterior articular surface: flat (0); concave (1); convex (2). The first caudal vertebra of *Patagotitan* has an unusual morphology having a flat anterior articulation surface and a markedly convex posterior one. Therefore, the character reflecting the first caudal centrum morphology (platycoelous, procoelous or biconvex) was split in two characters, one for the anterior articular surface and the second one for the posterior articular surface morphology. ***Bustingorrytitan=?***

226. First caudal neural arch, coel on lateral aspect of neural spine: absent (0); present (1). (Wilson, 2002:character 117) ***Bustingorrytitan=?***

227. Anterior caudal vertebrae (mainly the first and second): ventral bulge on transverse process: absent (0); present (1). (D'Emic, 2012:character 52) ***Bustingorrytitan=0***

228. Anterior caudal vertebrae, lateral pneumatic foramen: absent (0); present (1). The derived state is used even for small foramina. (Modified from D'Emic, 2012:character 56) ***Bustingorrytitan=0***

229. Posteriormost anterior and middle caudal vertebrae, transverse processes orientation: perpendicular (0); swept backwards, reaching the posterior margin of the centrum (1). (D'Emic, 2012:character 59)

Bustingorrytitan=?

230. Anterior caudal vertebrae, transverse processes: ventral surface directed laterally or slightly ventrally (0); directed dorsally (1). (Whitlock, 2011:character 125) ***Bustingorrytitan=?***

231. Anterior caudal centra (excluding the first), articular face shape: amphiplatyan or amphicoelous (0); procoelous/distoplatyan (1); slightly procoelous (2); procoelous (3); posterior surface markedly more concave than the anterior one (4). (Modified from González Riga *et al.*, 2009) **Bustingorrytitan=3**
232. Anterior caudal centra, large pneumatic foramen in lateral aspect (pleurocoels): absent or reduced (0); present (1). The derived state is solely used when the lateral pneumatic foramen is large, occupying much of the centrum, as those typically observed in several diplodocids. (Modified from Wilson, 2002:character 119)
- Bustingorrytitan=0**
233. Anterior caudal vertebrae, ventral surface: convex transversely (0); concave transversely (1). (Upchurch *et al.*, 2004:character 182) **Bustingorrytitan=1**
234. Anterior and middle caudal vertebrae, ventrolateral ridges: absent (0); present (1). (Upchurch *et al.*, 2004:character 183) **Bustingorrytitan=?**
235. Anterior and middle caudal vertebrae, triangular lateral process on the neural spine: absent (0); present (1). (Whitlock, 2011:character 123) **Bustingorrytitan=0**
236. Anterior caudal transverse processes shape: triangular, tapering distally (0); "wing like", not tapering distally (1); similar to the wing-shape process (not tapering distally), but with the high lateral margin dorsomedially oriented instead of vertical. (Modified from Wilson, 2002:character 128, adding one character for describing the high processes observed in some titanosauriforms.) **Bustingorrytitan=?**
237. Anterior caudal neural spines, transverse shape: anteroposteriorly long and lateromedially thin neural spines (0); square (length and width similar) (1); laterally expanded, around 1.5 times wider than long (2). (Modified from Wilson, 2002:character 126) **Bustingorrytitan=0**
238. Anterior caudal transverse processes, proximal depth: shallow, on centrum only (0); deep, extending from centrum to neural arch (1). (Wilson, 2002:character 127) **Bustingorrytitan=?**
239. Anterior caudal transverse processes, diapophyseal laminae (ACDL, PCDL, PRDL, PODL): absent (0); present (1). (Wilson, 2002:character 129) **Bustingorrytitan=?**
240. Anterior caudal transverse processes, anterior centrodiapophyseal lamina (ACDL), shape: single (0); divided (1). (Wilson, 2002:character 130) **Bustingorrytitan=?**
241. Anterior caudal vertebrae, hypophene ridge: absent (0); present (1). (Upchurch *et al.*, 2004:character 187)
- Bustingorrytitan=1**

242. Anterior caudal centra, length: approximately the same (0); or doubling over the first 20 vertebrae (1).
 (Wilson, 2002:character 120) **Bustingorrytitan=0**
243. Anterior caudal neural arches, spinoprezygapophyseal lamina (SPRL): absent, or present as small short ridges that rapidly fade out into the anterolateral margin of the spine (0); present, extending onto lateral aspect of neural spine (1); present, well developed and extending onto the anterior or anterolateral edges of the neural spine (2). (Modified from Wilson, 2002:character 121). A third state was incorporated in order to include the morphology observed in some taxa in which the SPRL is well developed, but is not extending into the lateral aspect of the neural spine, as is the case of *Patagotitan*. **Bustingorrytitan=?**
244. Anterior caudal neural arches, spinodiapophyseal lamina (SPDL): absent (0); present (1). In titanosaurs the SPDL, when present, is extending from the diapophyseal section of the transverse process (the dorsalmost part of it) up to the neural spine. **Bustingorrytitan=?**
245. Anterior caudal neural arches, spinoprezygapophyseal lamina (SPRL)- spinopostzygapophyseal lamina (SPOL) contact: absent (0); present, forming a prominent lamina on lateral aspect of neural spine (1).
 (Wilson, 2002:character 122) **Bustingorrytitan=0**
246. Anterior caudal neural arches, prespinal lamina (PRSL): absent (0); present (1). (Wilson, 2002:character 123) **Bustingorrytitan=?**
247. Anterior caudal vertebrae, ventral and medially placed SPRL, usually described as bifurcated PRSL: absent (0); present (1). This character was originally proposed as an autapomorphy of *Futalognkosaurus* but is not just restricted to this sauropod. **Bustingorrytitan=1**
248. Anterior caudal prespinal lamina (PRSL), triangular shaped product of a dorsal expansion of it: absent (0); present (1) **Bustingorrytitan=?**
249. Anterior caudal vertebrae, pair thin laminae that are bounding the prespinal laminae and that diverge dorsally: absent (0); present (1). This character was initially proposed as an autapomorphy of *Bonitasaura* but is present in some other sauropods, such as *Patagotitan*. **Bustingorrytitan=0**
250. Middle caudal centra, shape: cylindrical (0); with flat ventral margin (1); quadrangular, flat ventrally and laterally (2); trapezoidal (laterally compressed forming a shallow fossa) (3). (Modified from Wilson, 2002:character 131, Carballido *et al.*, 2020:character 250 and Hechenleitner *et al.*, 2020:character 250)
Bustingorrytitan=1

251. Anterior and middle caudal centra, ventral longitudinal hollow: absent (0); present (1). (Wilson, 2002:character 132) **Bustingorrytitan=1**
252. Middle caudal centra, articular face shape: amphiplatyan or amphicoelous (0); procoelous/distoplatyan (1); slightly procoelous (2); procoelous (3). (González Riga *et al.*, 2009) **Bustingorrytitan=3**
253. Posteriormost anterior and middle caudal vertebrae, location of the neural arches: over the midpoint of the centrum with approximately subequal amounts of the centrum exposed at either end (0); on the anterior half of the centrum (1). (Upchurch *et al.*, 2004:character 185) **Bustingorrytitan=1**
254. Anterior caudal vertebrae, anterior face of the centrum strongly inclined anteriorly: absent (0); present (1). (Santucci and Arruda Campos, 2011:character 256) **Bustingorrytitan=0**
255. Middle caudal vertebrae, with the anterior face strongly inclined anteriorly: absent (0); present (1)
Bustingorrytitan=0
256. Middle caudal vertebrae, height of the pedicels below the prezygapophysis: low with curved anterior edge of the pedicel (0); high with vertical anterior edge of the pedicel (1). (Carballido *et al.*, 2012)
Bustingorrytitan=0
257. Middle caudal vertebrae, orientation of the neural spines: anteriorly (0); vertical (1); slightly directed posteriorly (2); strongly directed posteriorly (3). (Modified from Wilson, 2002:character 133)
Bustingorrytitan=?
258. Posterior caudal vertebrae, neural spine strongly displaced posteriorly: absent (0); present (1). (Carballido *et al.*, 2012) **Bustingorrytitan=?**
259. Middle caudal vertebrae, ratio of centrum length to centrum height: less than 2, usually 1.5 or less (0); 2 or higher (1). (Upchurch *et al.*, 2004:character 179) **Bustingorrytitan=0**
260. Anterior-posterior caudal vertebrae (those with still well developed neural spine), neural spine orientation: vertical (0); slightly directed posteriorly (1); strongly directed posteriorly (2). (Carballido *et al.*, 2012)
Bustingorrytitan=?
261. Posterior caudal centra, articular face shape: amphyplatic (0); procoelous (1); opisthocoelous (2). (Modified from González Riga *et al.*, 2009) **Bustingorrytitan=1**
262. Posterior caudal centra, shape: cylindrical (0); dorsoventrally flattened, breadth at least twice height (1). (Wilson, 2002:character 135) **Bustingorrytitan=0**

263. Posterior caudal vertebrae, ratio of length to height: less than 5, usually 3 or less (0); 5 or higher (1).
 (Upchurch *et al.*, 2004:character 180) ***Bustingorrytitan=?***
264. Distalmost caudal centra, articular face shape: platycoelous (0); biconvex (1). (Wilson, 2002:character 136)
Bustingorrytitan=?
265. Distalmost biconvex caudal centra, number: 10 or fewer (0); more than 30 (1). (Wilson, 2002:character 137)
Bustingorrytitan=?
266. Distalmost biconvex caudal centra, length-to height ratio: less than 4 (0); greater than 5 (1). (Wilson, 2002:character 138) ***Bustingorrytitan=?***
267. Forked chevrons with anterior and posterior projections: absent (0); present (1). (Wilson, 2002:character 143) ***Bustingorrytitan=0***
268. Forked chevrons, distribution: distal tail only (0); throughout middle and posterior caudal vertebrae (1).
 (Wilson, 2002:character 144) ***Bustingorrytitan=?***
269. Chevrons, crus bridging dorsal margin of haemal canal: present (0); absent (1). (Wilson, 2002:character 145)
Bustingorrytitan=1
270. Chevron haemal canal, depth: short, approximately 25% (0); or long, approximately 50% chevron length (1).
 (Wilson, 2002:character 146) ***Bustingorrytitan=1***
271. Chevrons: persisting throughout at least 80% of tail (0); disappearing by caudal 30 (1). (Wilson, 2002:character 147) ***Bustingorrytitan=?***
272. Posterior chevrons, distal contact: fused (0); unfused (open) (1). (Wilson, 2002:character 148)
Bustingorrytitan=?

Shoulder girdle

273. Posture: bipedal (0); columnar, obligatory quadrupedal posture (1). (Wilson, 2002:character 149)
Bustingorrytitan=?

Scapular girdle

274. Scapular acromion process, size: Narrow (0); broad, width more than 150% minimum width of blade (1).
 (Wilson, 2002:character 150) ***Bustingorrytitan=1***

275. Scapular blade, orientation respect to coracoid articulation: perpendicular (0); forming a 45° angle (1). (Wilson, 2002:character 151) **Bustingorrytitan=1**
276. Scapular blade, distal expansion: absent (0); present (1) **Bustingorrytitan=?**
277. Scapular blade, shape: acromial edge not expanded (both edges are running parallel to each other) (0); rounded expansion on acromial side (1); racquet-shaped (2); marked distal expansion due to the posterodorsal orientation of the dorsal edge (3). (Wilson, 2002:character 152; as modified by Carballido *et al.*, 2017:character 277) **Bustingorrytitan=1**
278. Scapula, acromion process dorsal margin: concave or straight (0); with V-shaped concavity (1); with U-shaped concavity (2). (Sereno *et al.*, 2007:character 88) **Bustingorrytitan=?**
279. Scapula, highest point of the dorsal margin of the blade: lower than the dorsal margin of the proximal end (0); at the same height than the dorsal margin of the proximal end (1); higher than the dorsal margin of the proximal end (2). (Carballido *et al.*, 2012, from Mannion, 2009) **Bustingorrytitan=0**
280. Scapula, development of the acromion process: undeveloped (0); well developed (1). (Carballido *et al.*, 2012) **Bustingorrytitan=1**
281. Scapular length/minimum blade breadth: 5.5 or less (0); 5.5 or more (1). (Carballido *et al.*, 2012)
Bustingorrytitan=?
282. Scapula, ventral margin with a well-developed ventromedial process: absent (0); present, having only one (1); present, having two processes (2). (Modified from Carballido *et al.*, 2011) **Bustingorrytitan=0**
283. Scapular, acromial process position: lies nearly glenoid level (0); lies nearly midpoint scapular body (1). (Carballido *et al.*, 2012) **Bustingorrytitan=1**
284. Scapular acromion length: less than 1/2 scapular length (0); at least 1/2 scapular length (1). (Mannion *et al.*, 2012:character 168) **Bustingorrytitan=1**
285. Glenoid scapular orientation: relatively flat or laterally facing (0); strongly beveled medially (1). (Wilson, 2002:character 153) **Bustingorrytitan=0**
286. Scapular blade, cross-sectional shape at base: flat or rectangular (0); D-shaped (1). (Wilson, 2002:character 154) **Bustingorrytitan=?**
287. Coracoid, proximodistal length: less than the length of scapular articulation (0); approximately twice the length of scapular articulation (1). (Wilson, 2002:character 155) **Bustingorrytitan=?**

288. Coracoid, anteroventral margin shape: rounded (0); rectangular (1). (Wilson, 2002:character 156)

Bustingorrytitan=1

289. Dorsal margin of the coracoid in lateral view: reaches or surpasses the the level of the dorsal margin of the scapular expansion (0); lies below the level of the scapular proximal expansion and separated from the latter by a V-shaped notch (1). (Upchurch *et al.*, 2004:character 207) **Bustingorrytitan=1**

290. Coracoid, infraglenoid deep groove: absent (0); present (1) **Bustingorrytitan=1**

291. Coracoid, infraglenoid lip: absent (0); present (1). (Wilson, 2002:character 157) **Bustingorrytitan=1**

292. Sternal plate, shape: posterolateral margin curved (0); posterolateral margin expanded as a corner (1). (D'Emic, 2012:character 76) **Bustingorrytitan=?**

293. Sternal plate, shape: oval (0); crescentic (1). (Wilson, 2002:character 158) **Bustingorrytitan=1**

294. Prominent posterolateral expansion of the sternal plate producing a kidney-shaped profile in dorsal view: absent (0); present (1). (Upchurch *et al.*, 2004:character 211) **Bustingorrytitan=1**

295. Prominent parasagittal oriented ridge on the dorsal surface of the sternal plate: absent (0); present (1). (Upchurch *et al.*, 2004:character 212) **Bustingorrytitan=?**

296. Ridge on the ventral surface of the sternal plate: absent (0); present (1). (Upchurch *et al.*, 2004:character 213) **Bustingorrytitan=1**

297. Ratio of maximum length of sternal plate to the humerus length: less than 0.75, usually less than 0.65 (0); greater than 0.75 (1). (Upchurch *et al.*, 2004:character 209) **Bustingorrytitan=0**

Forelimbs

298. Humerus, strong posterolateral bulge around the level of the deltopectoral crest: absent (0); present (1). (D'Emic, 2012:character 80) **Bustingorrytitan=0**

299. Humerus, radial and ulnar condyles shape: radial condyle divided on anterior face by a notch (0); undivided (1). (D'Emic, 2012:character 83) **Bustingorrytitan=0**

300. Humerus-to-femur ratio: less than 0.60 (0); 0.60 to 0.69 (1); 0.70 to 0.90 (2); greater than 0.90 (3) (Modified from Upchurch *et al.*, 2004: character 216, Carballido *et al.*, 2017:character 300, and Hechenleitner *et al.*, 2020:character 300) **Bustingorrytitan=?**

301. Humeral deltopectoral attachment, development: prominent (0); reduced to a low crest or ridge (1). (Wilson, 2002:character 160) **Bustingorrytitan=1**
302. Humeral deltopectoral crest, shape: relatively narrow throughout length (0); markedly expanded distally (1). (Wilson, 2002:character 161) **Bustingorrytitan=1**
303. Humeral midshaft cross-section, shape: circular (0); elliptical (1). (Mannion *et al.*, 2011:character 170)
Bustingorrytitan=1
304. Humerus, RI (*sensu* Wilson and Upchurch, 2003): Gracile (less than 0.27) (0); medium (0.28-0.32) (1); Robust (more than 0.33) (2). (Carballido *et al.*, 2012) **Bustingorrytitan=0**
305. Humeral distal condyles, articular surface shape: restricted to distal portion of humerus (0); exposed on anterior portion of humeral shaft (1). (Wilson, 2002:character 163) **Bustingorrytitan=1**
306. Humeral distal condyle, shape: divided (0); flat (1). (Wilson, 2002:character 164) **Bustingorrytitan=1**
307. Humeral, lateral margin: medially deflected (0); almost straight until the half-length or even more (1); almost straight until the proximal third of the total length of the humerus (2). (Carballido *et al.*, 2012)
Bustingorrytitan=?
308. Humeral proximolateral corner, shape: rounded, the dorsal surface is well convex (0); pronounced / square, the dorsal surface low, almost flat (1). (Wilson, 2002:character 159) **Bustingorrytitan=1**
309. Ulnar proximal condyle, shape: subtriangular (0); triradiate, with deep radial fossa (1). (Wilson, 2002:character 165) **Bustingorrytitan=1**
310. Ulnar proximal condylar processes, relative lengths: subequal (0); unequal, anterior arm longer (1). (Wilson, 2002:character 166) **Bustingorrytitan=1**
311. Ulnar olecranon process, development: prominent, projecting above proximal articulation (0); rudimentary, level with proximal articulation (1). (Wilson, 2002:character 167) **Bustingorrytitan=1**
312. Ulna, length-to-proximal breadth ratio: gracile (0); stout (1). (Wilson, 2002:character 168)
Bustingorrytitan=1
313. Radial distal condyle, shape: round (0); subrectangular, flattened posteriorly and articulating in front of ulna (1). (Wilson, 2002:character 169) **Bustingorrytitan=1**
314. Radius, distal breadth: slightly larger than midshaft breadth (0); approximately twice midshaft breadth (1). (Wilson, 2002:character 170) **Bustingorrytitan=1**

315. Radius, distal condyle orientation: perpendicular to long axis of shaft (0); beveled approximately 20° proximolaterally relative to long axis of shaft (1). (Wilson, 2002:character 171) **Bustingorrytitan=1**
316. Carpal bones, number: 3 or more (0); 2 or fewer (1). (Wilson, 2002:character 173) **Bustingorrytitan=?**
317. Carpal bones, shape: round (0); block-shaped, with flattened proximal and distal surfaces (1). (Wilson, 2002:character 174) **Bustingorrytitan=?**
318. Metacarpus, shape: spreading (0); bound, with sub-parallel shafts and articular surfaces that extend half their length (1). (Wilson, 2002:character 175) **Bustingorrytitan=1**
319. Metacarpals, shape of proximal surface in articulation: gently curving, forming a 90 arc (0); U-shaped, subtending a 270 arc (1). (Wilson, 2002:character 176) **Bustingorrytitan=1**
320. Longest metacarpal-to-radius ratio: close to 0.3 (0); 0.45 or more (1). (Wilson, 2002:character 177)
Bustingorrytitan=1
321. Metacarpal I, length: shorter than metacarpal IV (0); longer than metacarpal IV (1). (Wilson, 2002:character 178) **Bustingorrytitan=1**
322. Metacarpal I, distal condyle shape: divided (0); undivided (1). (Wilson, 2002:character 179)
Bustingorrytitan=1
323. Metacarpal I distal condyle, transverse axis orientation: beveled approximately 20° respect to axis of shaft (0); proximodistally or perpendicular with respect to axis of shaft (1). (Wilson, 2002:character 180)
Bustingorrytitan=0
324. Manual digits II and III, phalangeal number: 2- 3-4-3-2 or more (0); reduced, 2-2-2- 2-2 or less (1); absent or unossified (2). (Wilson, 2002:character 181) **Bustingorrytitan=?**
325. Manual phalanx I.1, shape: rectangular (0); wedge-shaped (1). (Wilson, 2002:character 182)
Bustingorrytitan=?
326. Manual nonungual phalanges, shape: longer proximodistally than broad transversely (0); broader transversely than long proximodistally (1). (Wilson, 2002:character 183) **Bustingorrytitan=?**

Pelvic girdle

327. Pelvis, anterior breadth: narrow, ilia longer anteroposteriorly than distance separating preacetabular processes (0); broad, distance between preacetabular processes exceeds anteroposterior length of ilia (1). (Wilson, 2002:character 184) ***Bustingorrytitan=?***
328. Ilium, ischial peduncle size: large, prominent (0); low, rounded (1). (Wilson, 2002:character 185)
Bustingorrytitan=?
329. Ilium, dorsal margin shape: flat (0); semicircular (1). (Wilson, 2002:character 186) ***Bustingorrytitan=1***
330. Ilium, preacetabular ventral margin shape: straight (0), concave (1); with a convex ventral bump (2)
Bustingorrytitan=?
331. Ilium, preacetabular process shape: pointed, arching ventrally (0); semicircular, with posteroventral excursion of cartilage cap (1). (Wilson, 2002:character 188) ***Bustingorrytitan=1***
332. Ilium, preacetabular process orientation: anterolateral to body axis (0); perpendicular to body axis (1). (Wilson, 2002:character 189) ***Bustingorrytitan=?***
333. Highest point on the dorsal margin of the ilium: lies caudal to the base of the pubic process (0); lies cranial to the base of the pubic process (1). (Upchurch *et al.*, 2004:character 245) ***Bustingorrytitan=?***
334. Pubis length respect to ischium: pubis slightly smaller or subequal to ischium (0); pubis larger (120% +) than ischium (1). (Carballido *et al.*, 2012) ***Bustingorrytitan=?***
335. Pubis, ambiens process development: small, confluent with anterior margin of pubis prominent, (0); projects anteriorly from anterior margin of pubis (1). (Wilson, 2002:character 189) ***Bustingorrytitan=0***
336. Pubic apron, shape: flat (straight symphysis) (0); canted anteromedially (gentle shaped symphysis) (1). (Wilson, 2002:character 190) ***Bustingorrytitan=1***
337. Puboischial contact, length: approximately one third total length of pubis (0); one half total length of pubis (1). (Wilson, 2002:character 191) ***Bustingorrytitan=?***
338. Ischium, acetabular articular surface: maintains approximately the same transverse width throughout its length (0); is transversely narrower in its central portion and strongly expanded as it approaches the iliac and pubic articulations (1). (Mannion *et al.*, 2013:character 180) ***Bustingorrytitan=?***
339. Ischium, iliac peduncle with constriction or "neck": absent (0); present (1). (Whitlock, 2011:character 173)
Bustingorrytitan=?

340. Ischium, elongate muscle scar on proximal end: absent (0); present (1). (Whitlock, 2011:character 174)

Bustingorrytitan=?

341. Ischial blade, shape: emarginate distal to pubic peduncle (0); no emargination distal to pubic peduncle (1). (Wilson, 2002:character 193) ***Bustingorrytitan=?***

342. Ischia pubic articulation: less or equal to the anteroposterior length of pubic pedicel (0); greater than the anteroposterior length of pubic pedicel (1). (Salgado *et al.*, 1997) ***Bustingorrytitan=?***

343. Ischia, anteroposterior pubic pedicel width divided the total length of the ischium: less than 0.5 (0); 0.5 or larger (1). (Carballido *et al.*, 2012) ***Bustingorrytitan=?***

344. Ischial distal shaft, shape: triangular, depth of ischial shaft increases medially (0); bladelike, medial and lateral depths subequal (1). (Upchurch *et al.*, 2004:character 194) ***Bustingorrytitan=?***

345. Ischial distal shafts, cross-sectional shape: V-shaped, forming an angle of nearly 50° with each other (0); flat, nearly coplanar (1). (Wilson, 2002:character 195) ***Bustingorrytitan=?***

346. Ischia, distal end: is only slightly expanded (0); is strongly expanded dorsoventrally (1). (Upchurch, 1998:character 183) ***Bustingorrytitan=?***

347. Ischium, angle formed between the shaft and the acetabular line: forming an almost right angle (80-110°) (0) or; a close angle (less than 70°) (1). (Carballido *et al.*, 2012) ***Bustingorrytitan=?***

348. Ischial tuberosity: absent (0); present (1). The tuberosity noted by Otero (2010) for the ischium of *Neuquensaurus* (Otero, 2010, fig. 8) and is present in other taxa, such as *Patagotitan*, *Bonitasaura*, *Futalognkosaurus*, *Alamosaurus* and *Neuquensaurus*. ***Bustingorrytitan=?***

Hindlimbs

349. Femur, longitudinal ridge on the anterior face: absent (0); present (1). (D'Emic, 2012:character 107)

Bustingorrytitan=0

350. Femur, fibular condyle: well developed, having a similar posterior projection than the tibial one (0); much shorter than the tibial condyle (1). The fibular condyle of *Patagotitan* and *Bonitasaura* is reduced in its posterior projection respect to that of most other sauropods, which fibular and tibial condyles are almost equally posteriorly projected. ***Bustingorrytitan=0***

351. Femur, epicondyle development: well developed (0); reduced, almost absent (1). In *Patagotitan* the epicondyle is extremely developed and notorious in posterior and distal view, as a minor step laterally projected. In contrast, in some titanosaurs the epicondyle is almost imperceptible, as is the case of *Dreadnoughtus*, *Opsithocoelicaudia*, *Neuquensaurus*, and *Saltasaurus*. **Bustingorrytitan=1**
352. Femur, fourth trochanter position: almost at the half of the femur (0); in the proximal third of the femur (1). The fourth trochanter of *Patagotitan* is positioned around the proximal third of the total femur length, similar to the position observed in *Futalognkosaurus*, *Bonitasaura*, and some other non-Lognkosauria as *Rapetosaurus*, *Saltasaurus* and *Neuquensaurus*. In contrast, the fourth trochanter of most sauropods is around the half of the total femur length, being even some more lower in *Opisthocoelicaudia*. **Bustingorrytitan=0**
353. Femur, fourth trochanter development: prominent (0); reduced to crest or ridge (1); extremely reduced (2). (Modified from Wilson, 2002:character 196, following Whitlock, 2011:character 186) **Bustingorrytitan=1**
354. Femur, lesser trochanter: present (0); absent (1). (Wilson, 2002:character 197) **Bustingorrytitan=?**
355. Femur midshaft, transverse diameter: subequal to anteroposterior diameter (0); 125- 150% anteroposterior diameter (1); at least 185% anteroposterior diameter (2). (Wilson, 2002:character 198) **Bustingorrytitan=?**
356. Femur, lateral bulge (marked by the lateral expansion and a dorsomedial orientation of the laterodorsal margin of the femur, which starts below the femur head ventral margin): absent (0); present (1). (Salgado *et al.*, 1997) **Bustingorrytitan=1**
357. Femur, pronounced ridge on posterior surface between greater trochanter and head: absent (0); present (1). (Whitlock, 2011:character 181) **Bustingorrytitan=0**
358. Femur head position: perpendicular to the shaft, rises at the same level than the greater trochanter (0); dorsally directed, rises well above the level of the greater trochanter (1). (Modified from Upchurch *et al.*, 2004:character 263) **Bustingorrytitan=1**
359. Femur, distal condyles relative transverse breadth: subequal (0); tibial much broader than fibular (1). (Wilson, 2002:character 2000) **Bustingorrytitan=1**
360. Femur, distal condyles orientation: perpendicular or slightly beveled dorsolaterally (0); or beveled dorsomedially approximately 10 relative to femoral shaft (1). (Wilson, 2002:character 201)
Bustingorrytitan=1

361. Femur, distal condyles articular surface shape: restricted to distal portion of femur (0); expanded onto anterior portion of femoral shaft (1). (Wilson, 2002:character 202) **Bustingorrytitan=1**
362. Situation of the femoral fourth trochanter: on the caudal surface of the shaft, near the midline (0); on the caudomedial margin of the shaft (1). (Upchurch *et al.*, 2004:character 268) **Bustingorrytitan=1**
363. Tibial proximal condyle, shape: narrow, long axis anteroposterior (0); expanded transversely, condyle subcircular (1). (Wilson, 2002:character 203) **Bustingorrytitan=1**
364. Tibial cnemial crest, orientation: projecting anteriorly (0); or laterally (1). (Wilson, 2002:character 204)
Bustingorrytitan=?
365. Tibia, distal breadth: approximately 125% (0); more than twice midshaft breadth (1). (Wilson, 2002:character 205) **Bustingorrytitan=0**
366. Tibial distal posteroventral process, size: broad transversely, covering posterior fossa of astragalus (0); shortened transversely, posterior fossa of astragalus visible posteriorly (1). (Wilson, 2002:character 206)
Bustingorrytitan=?
367. Fibula, proximal tibial scar, development: not well-marked (0); well-marked and deepening anteriorly (1). (Wilson, 2002:character 207) **Bustingorrytitan=1**
368. Fibula, lateral trochanter: absent (0); present (1). (Wilson, 2002:character 208) **Bustingorrytitan=1**
369. Fibular distal condyle, size: subequal to shaft (0); expanded transversely, more than twice midshaft breadth (1). (Wilson, 2002:character 209) **Bustingorrytitan=?**
370. Fibular, proximal end, anterior crest: absent or poorly developed (0); well developed creating an interlocking proximal crus (1). (D'Emic, 2012:character 111) **Bustingorrytitan=0**
371. Fibula, shaft shape: straight, or slightly sigmoidal (0); sigmoid, such that the proximal and distal faces are angled relative to midshaft (1). (D'Emic, 2012:character 113) **Bustingorrytitan=0**
372. Astragalus, shape: at least 1.5 times wider than anteroposteriorly long (0); anteroposterior and transverse dimensions subequal (1). (D'Emic, 2012:character 115) **Bustingorrytitan=0**
373. Astragalus, shape: rectangular (0); wedge shaped, with reduced anteromedial corner (1). (Wilson, 2002:character 210) **Bustingorrytitan=1**
374. Astragalus, fibular facet: faces laterally (0); faces posterolaterally, anterior margin visible in posterior view (1). (Whitlock, 2011:character 186) **Bustingorrytitan=0**

375. Astragalus, foramina at base of ascending process: present (0); absent (1). (Wilson, 2002:character 211)

Bustingorrytitan=1

376. Astragalus, ascending process length: limited to anterior two-thirds of astragalus (0); extending to posterior margin of astragalus (1). (Wilson, 2002:character 212) **Bustingorrytitan=1**

377. Astragalus, posterior fossa shape: undivided (0); divided by vertical crest (1). (Wilson, 2002:character 213)

Bustingorrytitan=0

378. Astragalus, transverse length: 50% more than (0); or subequal to proximodistal height (1). (Wilson,

2002:character 214) **Bustingorrytitan=0**

379. Calcaneum: present (0); absent or unossified (1). (Wilson, 2002:character 215) **Bustingorrytitan=?**

380. Distal tarsals 3 and 4: present (0); absent or unossified (1). (Wilson, 2002:character 216) **Bustingorrytitan=?**

381. Metatarsus, posture: bound (0); spreading (1). (Wilson, 2002:character 217) **Bustingorrytitan=1**

382. Metatarsal I proximal condyle, transverse axis orientation: perpendicular to (0); angled ventromedially approximately 15° to axis of shaft (1). (Wilson, 2002:character 218) **Bustingorrytitan=1**

383. Metatarsal I distal condyle, transverse axis orientation: perpendicular to (0); angled dorsomedially to axis of shaft (1). (Wilson, 2002:character 219) **Bustingorrytitan=1**

384. Metatarsal III length divided by metatarsal I length: less than 1.3 (0); more than 1.3 (1). (González Riga *et al.*, 2016:character 331) **Bustingorrytitan=?**

385. Longest metatarsal: metatarsal III (0); metatarsal IV (1). (González Riga *et al.*, 2016:character 334)

Bustingorrytitan=?

386. Metatarsal I distal condyle, posterolateral projection: absent (0); present (1). (Wilson, 2002:character 220)

Bustingorrytitan=1

387. Metatarsal I, minimum shaft width: less than that of metatarsals II-IV (0); or greater than that of metatarsals III-IV (1). (Wilson, 2002:character 221) **Bustingorrytitan=1**

388. Metatarsal I and V proximal condyle, size: smaller than (0); or subequal to those of metatarsals II and IV (1). (Wilson, 2002:character 222) **Bustingorrytitan=?**

389. Metatarsal III length: more than 30% (0); or less than 25% that of tibia (1). (Wilson, 2002:character 223)

Bustingorrytitan=?

390. Metatarsals III and IV, minimum transverse shaft diameters: subequal to (0); or less than 65% that of metatarsals I or II (1). (Wilson, 2002:character 224) ***Bustingorrytitan=0***
391. Metatarsal IV, proximomedial end, shape: flat or slightly concave (0); possesses a distinct embayment (1). (D'Emic, 2012:character 117) ***Bustingorrytitan=?***
392. Metatarsal IV, distal end, orientation: roughly perpendicular to long axis of bone (0); beveled upwards medially (1). (D'Emic, 2012:character 118) ***Bustingorrytitan=?***
393. Metatarsal V, length: shorter than (0); or at least 70% length of metatarsal IV (1). (Wilson, 2002:character 225) ***Bustingorrytitan=0***
394. Pedal nonungual phalanges, shape: longer proximodistally than broad transversely (0); broader transversely than long proximodistally (1). (Wilson, 2002:character 226) ***Bustingorrytitan=?***
395. Pedal digits II-IV, penultimate phalanges, development: subequal in size to more proximal phalanges (0); rudimentary or absent (1). (Wilson, 2002:character 227) ***Bustingorrytitan=?***
396. Pedal unguals, orientation: aligned with (0); or deflected lateral to digit axis (1). (Wilson, 2002:character 228) ***Bustingorrytitan=?***
397. Pedal digit I ungual, length relative to pedal digit II ungual: subequal (0); 25% larger than that of digit II (1). (Wilson, 2002:character 229) ***Bustingorrytitan=0***
398. Pedal digit I ungual, length: shorter (0); or longer than metatarsal I (1). (Wilson, 2002:character 230) ***Bustingorrytitan=1***
399. Pedal ungual I, shape: broader transversely than dorsoventrally (0); sickle-shaped, much deeper dorsoventrally than broad transversely (1). (Wilson, 2002:character 231) ***Bustingorrytitan=1***
400. Pedal ungual II-III, shape: broader transversely than dorsoventrally (0); sickle shaped, much deeper dorsoventrally than broad transversely (1). (Wilson, 2002:character 232) ***Bustingorrytitan=1***
401. Pedal digit IV ungual, development: subequal in size to unguals of pedal digits II and III (0); rudimentary or absent (1). (Wilson, 2002:character 233) ***Bustingorrytitan=?***
402. Unguals of pedal digit II and III, proximal dimensions: as broad as deep (0); significantly broader than deep (1). (Allain and Aquesbi, 2008:character 253) ***Bustingorrytitan=?***
403. Number of phalanges in pedal digit II: 3 (0); 2 (1). (González Riga *et al.*, 2016:character 348) ***Bustingorrytitan=?***

404. Number of phalanges in pedal digit III: 4 (0); 3 (1). (González Riga *et al.*, 2016:character 349)

Bustingorrytitan=?

405. Number of phalanges in pedal digit IV: 3 or more (0); 2 (1); 1 (2). (González Riga *et al.*, 2016:character 350)

Bustingorrytitan=?

New characters introduced by Pérez Moreno *et al.* (2022) to the matrix of Gallina *et al.* (2021)

406. Postorbital, excluded from the infratemporal fenestra due to the articulation of the jugal with the squamosal: absent (0), present (1). (Canudo *et al.*, 2018) **Bustingorrytitan=?**

407. Squamosal, ventral shape: thin (0); broad (1). (Canudo *et al.*, 2018) **Bustingorrytitan=?**

408. Preantorbital fenestra development: small, differentiated from the posterior maxillary foramen in its direction (see Wilson and Sereno, 1998) (0); laterally opened middle sized fenestra (1); laterally opened large fenestra (2). (Canudo *et al.*, 2018) **Bustingorrytitan=?**

409. Mid- and posterior dorsal neural arches, centroprezygapophyseal fossa depth: shallow or absent (0); deep, passing nearly all the way through the neural arch. (Wilson and Allain, 2015:character 101)

Bustingorrytitan=0

410. Mid- Posterior dorsal vertebrae, parapophysis, position with respect to prezygapophyses: at the same level or below (0); well above (1). (Wilson and Allain, 2015:character 100) **Bustingorrytitan=0**

411. Posterior dorsal neural arches, centroprezygapophyseal lamina (CPRL), shape: single (0); divided (1). (Wilson and Allain, 2015:character 107) **Bustingorrytitan=?**

412. Posterior dorsal neural arches, spinoparapophyseal lamina (SPPL): absent (0); present (1). (Wilson and Allain, 2015:character 109) **Bustingorrytitan=?**

413. Middle caudal vertebrae, prezygapophyses orientation: anterodorsally oriented (around 45 degrees) (0); anteriorly oriented (nearly horizontal) (1). (Canudo *et al.*, 2018:character 413) **Bustingorrytitan=0**

414. Position of the single, ventral process (if 2 are present ambiguous score): below the acromion (0); on the scapular blade (posterior to the acromion) (1). This character was modified. Taxa without ventral process are scored as missing data (?). (Modified Carballido *et al.*, 2020:character 414, based on Mannion *et al.*, 2012)

Bustingorrytitan=?

415. Ilium, postacetabular posteroverentral edge: open concave (0); U-shaped notch (1); horizontal and low V-shaped notch (2). (Carballido *et al.*, 2020:character 415) **Bustingorrytitan=?**
416. Pubis, ischiatic articular surface: continuous without marked angle change (0); marked step formed by a proximal posterior directed surface and a more distal posterodorsal oriented surface (1). (Carballido *et al.*, 2020:character 416) **Bustingorrytitan=0**
417. Pubis, proximal symphysis: merges with the pubic shaft (0); forms a marked ventromedially directed process (1). (Carballido *et al.*, 2020:character 417) **Bustingorrytitan=1**
418. Anterior caudal vertebrae, deep and marked fossa on anterior aspect of the transverse process (prcdf): absent (0); present (1). **Bustingorrytitan=0**
419. Apex of the convexity of the posterior articulation on anterior and middle caudal vertebrae: concentrical or slightly displaced above the centrum midline (0); strongly displaced upward, so that the apex of the posterior articulation is flushed to the level of the dorsal margin of the centrum (1). (Santucci and Arruda Campos, 2011:character 235) **Bustingorrytitan=0**
420. Articular facets of the prezygapophyses on anterior and middle caudal vertebrae: normal, not expanded (0); wide, with a dorsal and a ventral expansion or protuberance (1). (Santucci and Arruda Campos, 2011:character 237) **Bustingorrytitan=0**
421. Prezygapophyses curved downward on anteriormost caudal vertebrae: absent (0); present (1) (Santucci and Arruda Campos, 2011:character 238). **Bustingorrytitan=?**
422. Postzygapophyses located on the anterior half of the centrum on anterior and middle caudal vertebrae: absent (0); present (1). (Santucci and Arruda Campos, 2011: character 239) **Bustingorrytitan=0**
423. Length proportions of the prezygapophyses with respect to the centrum length in middle caudal vertebrae: less than 40% (0); between 40-50% (1); more than 50% (2). (Santucci and Arruda Campos, 2011:character 241) **Bustingorrytitan=0**
424. Haemal arches with double articular facets set in a concave posterodorsal surface: absent (0); present (1). (Modified from Santucci and Arruda Campos, 2011:character 240). **Bustingorrytitan=0**
425. Haemal arches, distal portion of the articular process with lateral bulge: absent (0); present (1)
- Bustingorrytitan=0**

426. Haemal arches, articular area: with no well-developed posterior protuberance below (0); with well-developed posterior protuberance below (1) **Bustingorrytitan=0**
427. Anterior caudal vertebrae, neural arch, SPOL development: null (0); poorly developed causing the articular facet of the postzygapophysis to project slightly from the midline (1); strongly developed forming a bony process that supports the articular facet of the postzygapophysis (2) **Bustingorrytitan=1**
428. Middle caudal vertebrae, neural arch, SPOL development: null (0); poorly developed causing the articular facet of the postzygapophysis to project slightly from the midline (1); strongly developed forming a bony process that supports the articular facet of the postzygapophysis (2) **Bustingorrytitan=?**
429. Anterior caudal vertebrae, central, height difference between the anterior and posterior articular face: null (0); present but little developed (1); present and strongly developed (2) **Bustingorrytitan=0**
430. Middle caudal vertebrae, central, height difference between the anterior and posterior articular face: null (0); present but little developed (1); present and strongly developed (2) **Bustingorrytitan=0**
431. Posterior caudal vertebrae, central, height difference between the anterior and posterior articular face: null (0); present but little developed (1); present and strongly developed (2) **Bustingorrytitan=?**

Character scoring for *Bustingorrytitan shiva* gen. et sp. nov

??00100?????????????0?
 0?01?0????2000??01110-??????0?0????????????????1???0??111-
 ?????????????10????11011112?11??0--
 1011?1?????????00??00??301?0?0??10??0?1?01131000??0?10???0?11??11?1?01?0110??111?11?
 10
 00?111011?1111111?11110????1?1??01?????????00101??101?11?0?11?000101100??111?11??0?0?
 ??0111??????00?0?01000?00001?00?

SOM 2. Autapomorphic characters of *Bustingorrytitan shiva* gen. et sp. nov recovered in the analysis

Ventral surface of the cervical centrum concave transversely (123); absence of pleurocoels in cervical centrum (125); neural spine of middle to posterior dorsal vertebrae with subparallel lateral margins in anterior posterior view (159); prespinal lamina (prsl) rough and wide, extended through almost all the neural spine, in dorsal vertebrae (164); presence of a single lamina (the single tpol) supporting the hypophene or the postzygapophysis from below in mid and posterior dorsal vertebrae (181); pleurocoel with angular dorsal margin in middle to posterior dorsal vertebrae (187); dorsal margin of pleurocoels at the level of or higher than the dorsal margin of the centrum in posterior dorsal vertebrae (188); absence of lateral spinopostzygapophyseal lamina (lspol) in middle and posterior dorsal neural spine (200); height of neural arch below the postzygapophyses (pedicel) subequal to or greater than height of centrum in mid and posterior dorsal vertebrae (203); slightly opisthocoelous posterior dorsal vertebrae (206); solid, not pneumatized, neural arches in anterior caudal vertebrae (221); anterior caudal vertebrae with spinoprezygapophyseal laminae ventral and medially placed, usually described as bifurcated PRSL (246); middle caudal centra with flat ventral margin (249); well developed acromion process of scapula (279); acromial process placed at nearly midpoint of the scapular body (282); scapular acromion at least 1/2 of scapular length (283); glenoid scapular orientated relatively flat or laterally facing (284); dorsal margin of the coracoid lies, in lateral view, below the level of the scapular proximal expansion and separated from the latter by a V-shaped notch (288); radial condyle of humerus divided on anterior face by a notch (298); humerus gracile (RI less than 0,27) (303); ulnar olecranon process rudimentary (310); transverse axis of the distal condyle of metacarpal I beveled approximately 20° respect to axis of shaft (322); distal breadth of tibia approximately 125% its midshaft breadth (364); minimum transverse shaft diameters of metatarsals III and IV subequal to that of metatarsals I or II (389); metatarsal V shorter than metatarsal IV (392); mid caudal prezygapophyses anterodorsally oriented (around 45 degrees) (412); proximal symphysis of the pubis forming a marked ventromedially directed process (416); anterior caudal vertebrae with SPOL poorly developed causing the articular facet of the postzygapophysis to project slightly from the midline (426).

Bibliography

- Allain R, Aquesbi N. 2008. Anatomy and phylogenetic relationships of *Tazoudasaurus naimi* (Dinosauria, Sauropoda) from the late Early Jurassic of Morocco. *Geodiversitas* 30(2): 345–424.
- Canudo, J.I., Carballido, J.L., Garrido, A.C., and Salgado, L. 2018. A new rebbachisaurid sauropod from the Aptian–Albian, Lower Cretaceous Rayoso Formation, Neuquén, Argentina. *Acta Palaeontologica Polonica* 63(4): 679–691.
- Carballido, J.L., Pol, D., Cerda, I.A., and Salgado, L. 2011. The osteology of *Chubutisaurus insignis* del Corro, 1975 (Dinosauria, Neosauropoda) from the ‘Middle’ Cretaceous of central Patagonia, Argentina. *Journal of Vertebrate Paleontology* 31(1): 93–110.

- Carballido, J.L., Salgado, L., Pol, D., Canudo, J.I., and Garrido, A.C. 2012. A new basal rebbachisaurid (Sauropoda, Diplodocoidea) from the Early Cretaceous of the Neuquén Group; evolution and biogeography of the group. *Historical Biology* 24(6): 631–654.
- Carballido, J.L., Scheil, M., Knötschke, N., and Sander, P.M. 2020. The appendicular skeleton of the dwarf macronarian sauropod *Europasaurus holgeri* from the Late Jurassic of Germany and a re-evaluation of its systematic affinities. *Journal of Systematic Palaeontology* 18(9): 739–781.
- Curry Rogers, K.A. 2005. Titanosauria: a phylogenetic overview. In: Curry Rogers KA, Wilson JA (eds) *The sauropods: evolution and paleobiology*. University of California Press, Berkeley and Los Angeles, pp 50–103.
- D'Emic, M. D. 2012. The early evolution of titanosauriform sauropod dinosaurs. *Zoological Journal of the Linnean Society* 166(3): 624–671.
- Gallina, P.A., Canale, J.I., and Carballido, J.L. 2021. The earliest known titanosaur sauropod dinosaur. *Ameghiniana* 58 (1): 35–31 [10.1016/B978-0-323-60984-5.00062-7](https://doi.org/10.1016/B978-0-323-60984-5.00062-7)
- González Riga, B.J., Lamanna, M., Ortiz David, L. and Calvo, J.O. 2016. A gigantic new dinosaur from Argentina and the evolution of the sauropod hind foot. *Scientific Reports* 6:19165.
- González Riga, B.J., Ortíz David, L. 2014. A new titanosaur (Dinosauria, Sauropoda) from the Upper Cretaceous (Cerro Lisandro Formation) of Mendoza Province Argentina. *Ameghiniana* 51(1): 3–25.
- González Riga BJ, Previtera E, Pirrone CA (2009) *Malarguesaurus florenciae* gen. et sp. nov., a new titanosauriform (Dinosauria, Sauropoda) from the Upper Cretaceous of Mendoza, Argentina. *Cretaceous Research* 30(1): 135–148.
- Harris, J.D. 2006. The significance of *Suuwassea emiliae* (Dinosauria: Sauropoda) for flagellicaudatan intrarelationships and evolution. *Journal of Systematic Palaeontology* 4: 185–198.
- Hechenleitner, E.M., Leuzinger, L., Martinelli, A.G., Rocher, S., Fiorelli, L.E., Taborda, J.R.A., Salgado, L. 2020. Two Late Cretaceous sauropods reveal titanosaurian dispersal across South America. *Communications Biology* 3:622.
- Mannion, P.D., Upchurch, P., Mateus, O., Barnes, R., Jones, M.E.H. 2012. New information on the anatomy and systematic position of *Dinheirosaurus lourinhensis* (Sauropoda: Diplodocoidea) from the Late Jurassic of Portugal, with a review of European diplodocoids. *Journal of Systematic Palaeontology* 10: 521–551.
- Mannion, P.D. 2009. A rebbachisaurid sauropod from the Lower Cretaceous of the Isle of Wight, England. *Cretaceous Research* 30: 521–526.
- Mannion, P.D., Upchurch, P., Barnes, R.N., and Mateus, O. 2013. Osteology of the Late Jurassic Portuguese sauropod dinosaur *Lusotitan atalaiensis* (Macronaria) and the evolutionary history of basal titanosauriforms. *Zoological Journal of the Linnean Society* 168(1): 98–206.
- Mannion, P.D., Upchurch, P. and Hutt, S. 2011. New rebbachisaurid (Dinosauria: Sauropoda) material from the Wessex Formation (Barremian, Early Cretaceous), Isle of Wight, United Kingdom. *Cretaceous Research* 32 (6): 774–780.
- Pérez Moreno, A.N., Otero, A., Carballido, J.L., Salgado, L., and Calvo, J.O. 2022. The appendicular skeleton of *Rinconsaurus caudamirus* (Sauropoda: Titanosauria) from the Upper Cretaceous of Patagonia, Argentina. *Cretaceous Research* 142: 105389 <https://doi.org/10.1016/j.cretres.2022.105389>
- Pol, D., Garrido, A.C. and Cerda, I.A. 2011. A new sauropodomorph dinosaur from the Early Jurassic of Patagonia and the origin and evolution of the sauropod-type sacrum. *PLOS ONE*; 6(1):e14572.

- Rauhut, O.W.M., Carballido, J.L., and Pol, D. 2015. A diplodocid sauropod dinosaur from the Late Jurassic Cañadón Calcáreo Formation of Chubut, Argentina. *Journal of Vertebrate Paleontology* DOI: 10.1080/02724634.2015.982798.
- Remes, K., Ortega, F., Fierro, I., Joger, U., Kosma, R., Ferrer, J.M.M., Ide, O.A., and Maga, A.A. 2009. New basal sauropod dinosaur from the Middle Jurassic of Niger and the early evolution of Sauropoda. *PLoS One*; 4:e6924.1–13.
- Salgado, L., Coria, R.A., and Calvo, J.O. 1997. Evolution of titanosaurid sauropods. Part I. Phylogenetic Analysis based on the postcranial evidence. *Ameghiniana* 34(1): 3–32.
- Santucci, R.M., and Arruda-Campos, D.A.D. 2011. A new sauropod (Macronaria, Titanosauria) from the Adamantina Formation, Bauru Group, upper Cretaceous of Brazil and the phylogenetic relationships of Aeolosaurini. *Zootaxa* 3085(1): 1–33.
- Sereno, P.C., Wilson, J.A., Witmer, L.M., Whitlock, J.A., Maga, A., Ide, O., and Rowe, T.A. 2007. Structural extremes in a Cretaceous dinosaur. *PloS ONE* 2(11), e1230.
- Upchurch, P., Barrett, P.M., Galton, P.M. 2007. A phylogenetic analysis of basal sauropodomorph relationships: implications for the origin of sauropod dinosaurs. *Special Papers in Palaeontology* 77: 57–90.
- Upchurch, P. 1998. The phylogenetic relationships of sauropod dinosaurs. *Zoological Journal of the Linnean Society* 124(1): 43–103.
- Upchurch, P., Barrett, P.M., and Dodson, P. 2004. Sauropoda. In D. B. Weishampel, P. Dodson and H. Osmólska (Eds.), *The Dinosauria* (2nd ed., pp. 259–322). University of California Press.
- Whitlock, J.A. 2011. A phylogenetic analysis of Diplodocoidea (Saurischia: Sauropoda). *Zoological Journal of the Linnean Society* 161: 872–915.
- Wilson J.A. and Upchurch P. 2003. A revision of *Titanosaurus* Lydekker (Dinosauria-Sauropoda), the first dinosaur genus with a ‘Gondwanan’ distribution. *Journal of Systematic Palaeontology* 1: 125–160.
- Wilson, J.A., Sereno, P.C. 1998. Early evolution and higher-level phylogeny of sauropod dinosaurs. *Society of Vertebrate Paleontology Memoir* 5: 1–68.
- Wilson, J.A. 2002. Sauropod dinosaur phylogeny: critique and cladistic analysis. *Zoological Journal of the Linnean Society* 136(2): 215–275.
- Wilson, J.A. and Allain, R. 2015. Osteology of *Rebbachisaurus garasbae* Lavocat, 1954, a diplodocoid (Dinosauria, Sauropoda) from the early Late Cretaceous-aged Kem Kem beds of southeastern Morocco. *Journal of Vertebrate Paleontology* DOI: 10.1080/02724634.2014.1000701.
- Zaher, H., Pol, D., Carvalho, A.B., Nascimento, P.M., Riccomini, C., Larson, P., Juarez-Valieri, R., Pires-Domingues, R., da Silva, N. J., and Campos, D.A. 2011. A complete skull of an Early Cretaceous sauropod and the evolution of advanced titanosaurians. *PLoS ONE* 6(2), e16663.

Table 1. Measurements of the caudal vertebrae of *Bustingorrytitan shiva* gen. and sp. nov.

	p	ch	cw	cl	nah	vh
MMCH-Pv 59/4	anterior	18	15	21	22	40
MMCH-Pv 59/5	anterior	15	15	20	16 (ins)	36 (i)
MMCH-Pv 59/6	mid	13	11	14	-	-
MMCH-Pv 59/7	mid-post.	9.5	7.5	14	11 (ins)	20.5
MMCH-Pv 60/2	posterior	4	3	7	-	-

Abbreviations: position (p); centrum height (ch); centrum width (cw); total centrum length (cl); neural arch height (nah); vertebra height (vh). i: incomplete; ins: neural spine incomplete. Units in cm.

Table 2. Measurements of the humeri of *Bustingorrytitan shiva* gen. and sp. nov.

	p	l	pw	mw	mp	dw
MMCH-Pv 59/20	right	160	56	28	68	48
MMCH-Pv 59/21	left	170	56	26	68	48

Abbreviations: position (p); l (length); proximal width (pw); minimum width of the diaphysis (mw); minimum perimeter of the diaphysis (mp); distal width (dw). Units in cm.

Table 3. Measurements of the ulnae of *Bustingorrytitan shiva* gen. and sp. nov.

	p	pw	dw	mw	mp	l
MMCH-Pv 59/23	right	31 (d)	12	14	40	110
MMCH-Pv 59/24	left	35	15	12 (i)	39 (i)	110
MMCH-Pv 60/3	right	36	18	12	39	94

Abbreviations: position (p); proximal width (pw); distal width (dw); minimum width of the diaphysis (mw); minimum perimeter of the diaphysis (mp); length (l). d: deformed; i: incomplete. Units in cm.

Table 4. Measurements of the metacarpals of *Bustingorrytitan shiva* gen. and sp. nov.

	p	l	mp	l/mp	a	b	C	D
MMCH-Pv 59/25	I	51	27.5	1.85	19	10	17	10
MMCH-Pv 59/26	II	53	28.5	1.85	15.5	11.5	15	11
MMCH-Pv 59/27	III	53	26	2.03	14	13	15	9.5
MMCH-Pv 59/28	IV	46	24	1.91	19	7.5	14	10
MMCH-Pv 59/29	V	45.5	28.5	1.59	16	9	16	11
MMCH-Pv 60/4	III	45	22.5	2	14	14.5	15.5	7.5

Abbreviations: position (p); length (l); minimum perimeter (mp); length/minimum perimeter ratio (l/mp); greater axis of the proximal end (a); lesser axis of the proximal end (b); greater axis of the distal end (c); lesser axis of the distal end (d). Units in cm.

Table 5. Measurements of the femora of *Bustingorrytitan shiva* gen. and sp. nov.

	p	pw	dw	mw	mp	l
MMCH-Pv 59/30	right	57	-	34	76*	112 (i)
MMCH-Pv 60/5	right	-	46	28	66	124 (i)
MMCH-Pv 61/1	left	33 (d)	48 (d)	26 (d)	-	160 (d)
MMCH-Pv 62/1	right	-	33 (i)	-	-	23 (i)

Abbreviations: position (p); proximal width (pw); distal width (dw); minimum width of the diaphysis (mw); minimum perimeter of the diaphysis (mp); length (l). d: deformed; i: incomplete; *: inferred. Units in cm.

Table 6. Measurements of the tibiae of *Bustingorrytitan shiva* gen. and sp. nov.

	p	pw	dw	mw	mp	l
MMCH-Pv 59/31	right	40	-	19	50.5	66 (i)
MMCH-Pv 60/6	left	44	25	18	44	110
MMCH-Pv 62/2	left	-	29	-	-	55 (i)

Abbreviations: position (p); proximal width (pw); distal width (dw); minimum width of the diaphysis (mw); minimum perimeter of the diaphysis (mp); length (l); i: incomplete. Units in cm.

Table 7. Measurements of the metatarsals of *Bustingorrytitan shiva* gen. and sp. nov.

	p	l	mw	mp	1/mp	a	b	c	d
MMCH-Pv 59/35	I left	16	9	30	0.53	16	10	7.5	13
MMCH-Pv 59/36	IV?	27 (i)	7	27	-	-	-	-	-
MMCH-Pv 59/37	V right	24	8	28	0.85	17	7.5	9	8.5

Abbreviations: position (p); length (l); minimum width of the diaphysis (mw); minimum perimeter of the diaphysis (mp); length/minimum perimeter of the diaphysis ratio (L/mp); greater axis of the proximal end (a); lesser axis of the proximal end (b); greater axis of the distal end (c); lesser axis of the distal end (d). i: incomplete. Units in cm.

Table 8. Measurements of the ungual phalanges of *Bustingorrytitan shiva* gen. and sp. nov.

	p	l	h	w
MMCH-Pv 59/38	I	17	10.5	7
MMCH-Pv 59/39	II	16	10	6
MMCH-Pv 59/40	III	14.5	8	6

Abbreviations: position (p); length (l); height (h); width (w). Units in cm.