

Middle Devonian and Frasnian bryozoan fauna from the Holy Cross Mountains, Poland

ANDREJ ERNST, GRZEGORZ RACKI, and PATRICK N. WYSE JACKSON



Ernst, A., Racki, G., and Wyse Jackson, P.N. 2026. Middle Devonian and Frasnian bryozoan fauna from the Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* 71 (1): 1–17.

The present paper describes a bryozoan fauna from the Devonian of Holy Cross Mountains (Poland) which includes nine species. Six species were described from the Dobruchna Member of the Skały Formation (Eifelian) of Skały section: *Cyclotrypa* sp., *Fistuliphragma gracilis*, *Leioclema passitabulatum*, *Intrapora leunissenii*, *Laxifenestella* sp., *Hemitrypella nodulosa*. The cryptostome species *Bigeyella indigena* was identified from the Laskowa Góra Beds (upper Givetian) of Józefka Hill at Górnó. The fenestrate *Rectifenestella* sp. is described from the lower Frasnian Kowala Formation (Kadzielnia Member) of Bolechowice and Kielce. The new monotypic fenestrate genus *Juanopora elegans* gen. et sp. nov. is described from the Wietrznia Beds (upper Givetian) of Wietrznia-I quarry at Kielce. The described bryozoans show palaeobiogeographical connections to the Middle–Upper Devonian of Germany, Spain, and Armenia. The bryozoan faunas of the Holy Cross Mountains are in accordance with the worldwide acme of the sessile filter-feeding benthos during the Givetian age, followed by a two-stage decline in diversity due to the Taghanic and Frasnian crises.

Key words: Bryozoa, taxonomy, evolution, palaeobiogeography, Devonian, Poland.

Andrej Ernst [Andrej.Ernst@uni-hamburg.de; ORCID: <https://orcid.org/0000-0002-1299-2559>], Institut für Geologie, Universität Hamburg, D-20146 Hamburg, Germany.

Grzegorz Racki [grzegorz.racki@us.edu.pl; ORCID: <https://orcid.org/0000-0003-4609-8341>], Institute of Earth Sciences, Department of Natural Sciences, University of Silesia in Katowice, Będzińska str. 58, PL 41-200 Sosnowiec, Poland.

Patrick N. Wyse Jackson [wysjcknp@tcd.ie; ORCID: <https://orcid.org/0000-0001-5605-0670>], Department of Geology, Trinity College, Dublin 2, Ireland.

Received 11 August 2025, accepted 18 November 2025, published online 19 February 2026.

Copyright © 2026 A. Ernst et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

In the Middle to Upper Devonian carbonate succession of the Holy Cross Mountains in south-central Poland (Fig. 1A), bryozoans are mostly a minor component in numerous benthic assemblages, which are usually dominated by brachiopods. Although reported in several publications (e.g., Pajchłowa 1957; Racki et al. 1985; Racki 2024), since the monographic work of Gürich (1896) this fossil group has been taxonomically studied exclusively from open shelf facies of the northern Łysogóry Region by Kiepurka (1965, 1973). Research was later significantly extended to the southern (Kielce) carbonate platform, and included many localities representing lithology variations from clayey-marly strata to stromatoporoid-coral reef-type limestones (Morozova et al. 2002).

The present contribution is a continuation of the work of Morozova et al. (2002), based on new specimens located in museum collections and on samples more recently collected during fieldwork undertaken by the second author and

students of the University of Silesia. Its main objective is to provide a taxonomic analysis of bryozoan material from five Middle Devonian and Frasnian sites located in the north-western part of the Holy Cross Mountains (Figs. 1B and 2), including providing important data from the famous Eifelian “brachiopod shale” of the Skały Formation (Pajchłowa 1957; Racki et al. 2022). The nine taxa described here include a new monotypic genus of fenestrellids. The migration history of the bryozoan faunas during gradual drowning of the Kielce carbonate platform (Fig. 2) is updated, as is information pertaining to their regional succession and zoogeographical affinities with Western Europe, especially the Eifel Mountains.

Institutional abbreviations.—GIUS, Institute of Earth Sciences of the University of Silesia in Katowice, Sosnowiec, Poland.

Nomenclatural acts.—This published work and the nomenclatural acts it contains, have been registered in Zoobank: urn:lsid:zoobank.org:pub:80C55D6F-6D65-40A6-AAE6-D26510271F92

Material and methods

The studied material comprises five samples containing loose fragments of bryozoan colonies. The friable marly shale and compact limestone samples come from following localities in Holy Cross Mts, Poland (Fig. 1B): Skały; Górno, Józefka Hill; Kielce, Wietrznia-I quarry; Bolechowice, Łgawa Hil; and Kielce, Cmentarna Hill (for details, see the section on the geological setting).

Thin sections were made from the studied samples. The separate fragments were imbedded in epoxide resin (SpeciFix-20), and then cut and polished. In total, 46 thin sections were made (24 × 48 mm). The studied material is deposited at the Institute of Earth Sciences of the University of Silesia in Katowice, Sosnowiec, Poland.

Bryozoan morphology has been studied using a binocular microscope. Morphologic character terminology is partly adopted from Boardman (1960) and Anstey and Perry (1970) for trepostomes, from Hageman (1991a, b) and Snyder (1991) for fenestrates. The following morphologic characters were measured and used for statistics in the studied material: branch width, exo- (endo-) zone width, axial ratio (ratio of endozone width to the branch width), autozoecial aperture width, aperture spacing (along branch, diagonally), acanthostyle diameter, exilazooecia or metazooecia width, wall thickness in exozone, vesicle diameter, number of vesicles per aperture, dissepiment width, fenestrule width

(length), distance between branch (dissepiment) centres, number of apertures per fenestrule length, maximal chamber width, width of superstructure openings. The spacing of structures is measured as a distance between their centres. Statistics were summarized using arithmetic mean, sample standard deviation, coefficient of variation, and minimum and maximum values.

Geological setting

The Middle to Upper Devonian epicontinental succession in the Holy Cross Mountains records a continuous stepwise drowning of an increasingly differentiated, vast bank-to-reef platform (Racki 1993; Szulczewski 1995; Narkiewicz et al. 2006; Racki et al. 2022), in general agreement with the transgressive-regressive cyclicity of the Euramerican eustatic curve of Johnson et al. (1985; Fig. 2).

The Holy Cross Mountains area is divided into two distinct paleogeographic-tectonic domains: the Kielce Paleohigh and the Łysogóry Paleolow, corresponding to a carbonate platform and a hemipelagic basin, respectively (Fig. 1B). The major facies regions are coupled with the transitional Kostomłoty Facies Zone (Racki 1993), which provides an opportunity to compare the bryozoan record in different sedimentary regimes (Fig. 2). However, the subsymmetric facies plan in the Late Devonian is shown by the central lo-




Fig. 1. Location of bryozoan sites in Poland (A) and the Holy Cross Mountains (B, based on Racki 1993: fig. 2) against paleogeography and regional geology; see Fig. 2 for refined stratigraphic-facies setting.




Fig. 2. Position of the bryozoan-bearing sections under this study (see Fig. 1B) relative to the developmental stages of the Middle–Late Devonian bank-to-reef complex of the Holy Cross Mountains; idealized stratigraphic-facies cross section according to Racki (1993: fig. 3, modified) to emphasize eustatic control of the rhythmic depositional pattern and major bryozoan immigration events; transgressive-regressive cycles Ic–IId after Johnson et al. (1985, modified; see Halamski et al. 2022: fig. 7); selected global biotic events following Racki (1993) and Becker (2025).

cation of the Dyminy Reef, surrounded by intrashelf basins: localized Chęciny-Zbrza (southern) and larger Łysogóry-Kostomłoty (northern), as summarized in Racki (1993), Szulczewski (1995) and Racki et al. (2022).

Three of the studied outcrops have already been investigated by Morozova et al. (2002, 2006). The most important data are provided for the uppermost Eifelian (*Polygnathus ensensis* Conodont Zone) “brachiopod shales” in the Łysogóry Region (Figs. 1B and 2), recently formalized as the Dobručna Member in the basal part of the shaly-calcareous Skały Formation (Racki et al. 2022). Two studied samples of fossiliferous marls were collected by graduate students of the University of Silesia from trench A dug in 1989 (Czaplikowska 1990; Woźniak 1992; see Woźniak et al. 2022).

Some species come from two upper Givetian localities in the northern part of the Kielce Region: the Górnio-Józefka hill (road cut, set A) and the Wietrznia-I quarry (section

Wi-IA, basal set A). They represent the bryozoan-enriched calcareous-marly succession of the Laskowa Góra Beds and their lateral equivalents in the basal Wietrznia Beds, roughly dated as an undivided interval of the *Schmidtognathus hermanni* to *Mesotaxis falsovalis* conodont zones (Makowski in Racki et al. 1993; Racki and Bultynck 1993; Baliński et al. 2016).

The relatively frequently encountered Frasnian fenestrellids in the northern foreslope successions of the Dyminy Reef, especially in the Kadzielnia Limestone Member of the Kowala Formation, have been recorded since the 19th century (see Racki 2024). They are described here for the first time from two localities of microbial-metazoan mud-mounds, Cmentarna Hill (set B1; Gawlik in Racki et al. 1993) in Kielce and the Łgawa Hill (set J; Racki 1993) near Bolechowice. The fossiliferous mid-slope bioconstructions are uncertainly dated as *Palmatolepis transitions* to *Palmatolepis punctata* conodont zones (Racki 2024).

Systematic palaeontology

Phylum Bryozoa Ehrenberg, 1831

Class Stenolaemata Borg, 1926

Superorder Palaeostomata Ma et al., 2014

Order Cystoporata Astrova, 1964

Suborder Fistuliporina Astrova, 1964

Family Fistuliporidae Ulrich, 1882

Genus *Cyclotrypa* Ulrich, 1896

Type species: Fistulipora communis Ulrich, 1890; Middle Devonian; Iowa, USA.

Emended diagnosis.—Encrusting colonies, often with multiple overgrowths. Cylindrical autozooezia with thin walls and complete diaphragms. Autozooezial apertures rounded. Lunaria absent. Autozooezia separated by extrazoidial vesicular skeleton. Acanthostyles occurring in roofs of vesicles. Low maculae often developed.

Remarks.—*Cyclotrypa* Ulrich, 1896, differs from *Fistulipora* M'Coy, 1849, and *Eridopora* Ulrich, 1882, in the absence of lunaria.

Stratigraphic and geographic range.—Silurian to Permian; Europe, North America, Asia.

Cyclotrypa sp.

Fig. 3A, B; Table 1.

Material.—Single colony, two thin sections (GIUS 4-2239 Sk/Bry-XIV-5N-1, -5N-2) from Dobruchna Member in the bottom part of the Skąły Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone); Skąły (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Description.—Encrusting colony, 1.75 mm in thickness. Autozooezia growing from thin epitheca, bending sharply at their bases towards colony surface. Autozooezial aper-

Table 1. Summary of descriptive statistics for *Cyclotrypa* sp. (single colony measured). Abbreviations: CV, coefficient of variation; MAX, maximal value; MIN, minimal value; N, number of measurements; SD, sample standard deviation; X, mean.

	N	X	MIN	MAX	SD	CV
Aperture width (mm)	20	0.30	0.27	0.34	0.021	7.14
Aperture spacing (mm)	20	0.49	0.43	0.55	0.030	6.07
Vesicle width (mm)	20	0.16	0.11	0.25	0.038	23.36
Vesicles per aperture	7	7.0	6.0	9.0	1.155	16.50

Table 2. Summary of descriptive statistics for *Fistuliphragma gracilis* Ernst, 2008 (five colonies measured). Abbreviations: CV, coefficient of variation; MAX, maximal value; MIN, minimal value; N, number of measurements; SD, sample standard deviation; X, mean.

	N	X	MIN	MAX	SD	CV
Aperture width (mm)	25	0.17	0.13	0.21	0.023	13.77
Aperture spacing (mm)	25	0.39	0.32	0.50	0.042	10.92
Vesicle width (mm)	25	0.09	0.06	0.14	0.023	24.00

tures circular to oval. Basal diaphragms absent. Vesicles large, separating autozooezia in 1–2 rows, 6–9 surrounding each autozooezial aperture, polygonal in tangential section, box-like to hemispheric, with plane or concave roofs, consisting of granular skeleton. Autozooezial walls thick, laminated.

Remarks.—*Cyclotrypa* sp. is similar to *Cyclotrypa communis* (Ulrich, 1890), from the Lower to Middle Devonian (Emsian–Givetian) of USA and Europe, but differs in having larger autozooezial apertures (average aperture width 0.30 mm vs. 0.20 mm in *C. communis*). *Cyclotrypa* sp. is similar to *C. cyclostoma* (Schlütter, 1889) [= *C. nekhoro-shevi* Kiepara, 1973] from the Middle Devonian (Eifelian–Givetian) of Europe and North Africa, but differs in the absence of apertural nodes.

Genus *Fistuliphragma* Bassler, 1934

Type species: Fistulipora spinulifera Rominger, 1866; Traverse Group, Middle Devonian; Michigan, USA.

Emended diagnosis.—Solid ramose and encrusting colonies, the latter usually in form of hollow tubes. Secondary overgrowth occurring. Autozooezia growing from a thin epitheca, bending sharply at their bases towards colony surface, with circular apertures and large, prominent lunaria. Hemiphragms present, positioned in alternating pattern in autozooezia, originating in laminated skeleton of autozooezia, rare in endozones, rare to abundant in exozones. Basal diaphragms rare to abundant. Vesicular skeleton well-developed. In the centre of each vesicle roof a single acanthostyle is present. Monticules present.

Remarks.—*Fistuliphragma* Bassler, 1934, is similar to *Cliotrypa* Ulrich & Bassler in Bassler, 1929, and *Strotopora* Ulrich in Miller, 1889, in the presence of hemiphragms, but differs from both genera in the absence of gonozooezia.

The species *Fistuliphragma crustiformis* Yang & Lu, 1983 (originally “*crustiforme*”, changed herewith to “*crustiformis*”, according to the Article 11.9 of the Code of Zoological Nomenclature), reported from the Pennsylvanian (Kasimovian–Gzhelian) of China (Xinjiang) belongs to the genus *Xiapora* Ernst & Gorgij, 2013. This species possesses incomplete vesicles which were interpreted as hemiphragms (Yang and Lu 1983: 271, pl. 5: 8–10).

Stratigraphic and geographic range.—Lower–Middle Devonian (Emsian–Givetian); Europe, China, North Africa, North America.

Fistuliphragma gracilis Ernst, 2008

Fig. 3C, D; Table 2.

2008 *Fistuliphragma gracilis* sp. nov.; Ernst 2008: 329–332, fig. 11A–G.

2011 *Fistuliphragma gracilis* Ernst, 2008; Ernst et al. 2011: 305–307, fig. 3e–g.

2012 *Fistuliphragma gracilis* Ernst, 2008; Ernst et al. 2012: 741–742, fig. 7a–d.

Material.—Five colonies, nine thin sections (GIUS 4-2239




Fig. 3. Cystoporate and trepostome bryozoans from the Devonian of Poland, Dobruchna Member in the bottom part of the Skaly Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone), Skaly. **A, B.** *Cyclotrypa* sp. **A.** Longitudinal section showing autozooezia and vesicles (GIUS 4-2239 Sk/Bry-XIV-5N-1) (A₁, A₂). **B.** Tangential section showing autozooezial apertures and vesicles (GIUS 4-2239 Sk/Bry-XIV-5N-2) (B₁, B₂). **C, D.** *Fistuliphragma gracilis* Ernst, 2008. **C.** Longitudinal section of a tubular colony showing autozooezial chamber, vesicles and hemiphragms (GIUS 4-2239 Sk/Bry-XIV-5A-2) (C₁–C₃). **D.** Tangential section showing autozooezial apertures with lunaria and vesicles (GIUS 4-2239 Sk/Bry-XIV-5E-1) (D₁, D₂). **E, F.** *Leioclema passitabulatum* Duncan, 1939. **E.** Longitudinal section showing autozooezial chambers and mesozooezia (GIUS 4-2239 Sk/Bry-XIV-5L-1). **F.** Tangential section showing autozooezial apertures, mesozooezia and acanthostyles (GIUS 4-2239 Sk/Bry-XIV-5H-1) (F₁, F₂).

Sk/Bry-XIV-5A-1, -5A-2, -5D, -5E-1, -5E-2, -5G-1, -5G-2, -5O-1, 5O-2) from Dobruchna Member in the bottom part of the Skaly Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone); Skaly (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Description.—Encrusting or hollow ramose colonies formed by encrusting overgrowths of ephemeral cylindrical objects. Encrusting sheets 0.6–1.0 mm thick. Autozooezia moderately long, narrow in endozones. Diaphragms absent. Hemiphragms abundant, long, proximally curved;

alternating in outer endozone and in exozone. Vesicles in outer endozone and exozone, 7–9 surrounding each autozoocial aperture, polygonal in tangential section, box-like to hemispheric, with plane or concave roofs, consisting of granular skeleton. Vesicular roofs containing acanthostyles, 0.018–0.025 mm in diameter. Autozoocial walls granular, 0.005–0.010 mm thick in endozones; laminated, 0.010–0.015 mm thick in exozones. Lunaria in outer exozone and endozone, horse-shoe shaped, large, with thick skeletal deposits. Maculae not observed.

Remarks.—*Fistuliphragma gracilis* Ernst, 2008, differs from *Fistuliphragma eifelensis* Ernst, 2008, in having smaller colonies, and smaller autozoocia (average autozoocial aperture width 0.17 mm vs. 0.28 mm in *F. eifelensis*).

Stratigraphic and geographic range.—Middle Devonian (lower Eifelian), Kierspe, Germany. Middle Devonian (upper Eifelian), western Rhenish Massif, Germany. Santa Lucia Formation, Lower–Middle Devonian (upper Emsian–lower Eifelian), Abeltas, Paradilla, Cantabrian Mountains, NW Spain. Dobruchna Member in the bottom part of the Skały Formation, Middle Devonian (uppermost Eifelian, *Polygnathus ensensis* Conodont Zone), Skały (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Order Trepostomata Ulrich, 1882

Suborder Halloporina Astrova, 1965

Family Heterotrypidae Ulrich, 1890

Genus *Leioclema* Ulrich, 1882

[= *Lioclema* Ulrich, 1882]

Type species: *Callopora punctata* Hall, 1858; Mississippian; Iowa, USA.

Emended diagnosis.—Encrusting, branched, less commonly massive colonies. Autozoocia with polygonal to rounded-polygonal, sometimes petaloid apertures. Autozoocial diaphragms rare. Mesozoocia abundant, with abundant diaphragms, often beaded. Acanthostyles abundant, commonly large. Autozoocial walls thin in endozone; laminated, regularly thickened in exozones (modified after Astrova 1978).

Remarks.—*Leioclema* Ulrich, 1882, differs from *Heterotrypa* Nicholson, 1879, in having rare autozoocial diaphragms and abundant acanthostyles and mesozoocia, and from *Stigmatella* Ulrich & Bassler, 1904, in having abundant mesozoocia.

Table 3. Summary of descriptive statistics for *Leioclema passitabulatum* Duncan, 1939 (three colonies measured). Abbreviations: CV, coefficient of variation; MAX, maximal value; MIN, minimal value; N, number of measurements; SD, sample standard deviation; X, mean.

	N	X	MIN	MAX	SD	CV
Aperture width (mm)	35	0.18	0.13	0.23	0.024	13.31
Aperture spacing (mm)	34	0.34	0.25	0.43	0.050	14.66
Mesozoecium width (mm)	34	0.09	0.04	0.14	0.022	25.62
Acanthostyle diameter (mm)	17	0.031	0.025	0.040	0.005	14.68
Mesozoocia per aperture	16	8.6	6.0	11.0	1.147	13.30

Stratigraphic and geographic range.—Lower Silurian to Pennsylvanian (Carboniferous); worldwide.

Leioclema passitabulatum Duncan, 1939

Fig. 3E–F; Table 3.

1939 *Leioclema passitabulatum* sp. nov.; Duncan 1939: 251, pl. 16: 8–10.

1964 *Leioclema passitabulatum* Duncan, 1939; Astrova 1964: 33, pl. 10: 2a, b.

2007 *Leioclema passitabulatum* Duncan, 1939; Ernst and Schroeder 2007: 210, figs. 2I–K, 3A–C.

2009 *Leioclema passitabulatum* Duncan, 1939; Ernst and May 2009: 779, figs. 7.1–7.4.

2012 *Leioclema passitabulatum* Duncan, 1939; Ernst et al. 2012: 743, fig. 8a–d.

Material.—Three colonies, three thin sections (GIUS 4-2239 Sk/Bry-XIV-5H-1, -5J-1, -5L-1) from Dobruchna Member in the bottom part of the Skały Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone); Skały (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Description.—Encrusting colonies, 0.90–1.08 mm in thickness. Autozoocia budding from a thin epitheca, briefly oriented parallel to the substrate, then bending sharply and intersecting the colony surface at right angles. Epitheca 0.013–0.018 mm thick. Autozoocial apertures rounded-polygonal to petaloid due to indenting acanthostyles. Autozoocial diaphragms few to absent, thin, straight or slightly deflected proximally. Mesozoocia abundant, 6–11 surrounding each aperture, polygonal in cross section, slightly beaded, containing planar diaphragms. Acanthostyles small, abundant, 1–5 surrounding each aperture, originating from the base of exozone, often indenting autozoocia, having distinct calcite cores and dark laminated sheaths. Autozoocial walls granular, in endozone 0.010–0.015 mm thick; in exozone 0.025–0.030 mm thick, distinctly laminated. Maculae not observed.

Remarks.—*Leioclema passitabulatum* Duncan, 1939, is similar to *Leioclema alpenense* Duncan, 1939, but differs from it in containing more abundant acanthostyles (1–5 vs. 2 acanthostyles around each aperture, respectively).

Stratigraphic and geographic range.—Lower Devonian, Lochkovian, Bortschovski Horizon, Podolia, Ukraine. Lower Devonian, Pragian, Koněprusy Limestone, Czech Republic. Middle Devonian, Eifelian, Traverse Group, Michigan, USA. Middle Devonian, lower Eifelian, Kierspe, Germany. Dobruchna Member in the bottom part of the Skały Formation, Middle Devonian (uppermost Eifelian (*Polygnathus ensensis* Conodont Zone), Skały (trench A dug in 1989; see Woźniak et al. 2022), Poland. Cürten Formation, Middle Devonian, lower Givetian, Dollendorf Syncline, Rhenish Slate Massif, Germany.

Order Cryptostomata Vine, 1884

Suborder Rhabdomesina Astrova & Morozova, 1956

Family Lenaporidae Ernst & Königshof, 2010




Fig. 4. Cryptostome and fenestrate bryozoans from the Devonian of Poland. A–C. *Bigeyella indigena* (Morozova & Weiss in Morozova et al., 2002), from the Laskowa Góra Beds, Middle Devonian (upper Givetian) (?*Polygnathus ansatus*–*Schmidtognathus hermanni*, and *Klapperina disparilis* conodont zones) of Gómo, Józefka Hill. A. Transverse section (GIUS 4-292 Gó/Bry A-8a). B. Longitudinal section (GIUS 4-292 Gó/Bry A-4b). C. Tangential section (GIUS 4-292 Gó/Bry A-3a) (C₁, C₂). D, E. *Intrapora leuisseni* Ernst, 2008, from the Dobruchna Member in the bottom part of the Skaly Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone) of Skaly. D. Transverse section showing autozoecia, metazoecia, and mesotheca {?} (GIUS 4-2239 Sk/Bry-XIV-5C-2) (D₁, D₂). E. Tangential section showing autozoecia, metazoecia, and acanthostyles (GIUS 4-2239 Sk/Bry-XIV-5I-1). F, G. *Rectifenestella* sp. from the Kadzielnia Limestone Member of the Kowala Formation, Lower Frasnian (*Palmatolepis transitans* Conodont Zone). F. Tangential section showing autozoecial chambers (GIUS 4-259 GC/Bry-1a), Bolechowice, Łgawa Hill (F₁, F₂). G. Apertures with stellate structures (arrows) (GIUS 4-256 GL/Bry-1b), Kielce, Cmentarna Hill.

Genus *Bigeyella* Morozova & Weiss in Morozova et al., 2006

Type species: Bigeyella sparsa Morozova & Weiss in Morozova et al., 2006; Middle Wietrznia Series (Member C), Upper Devonian (Frasnian); central Poland.

Emended diagnosis.—Branched colonies. Autozoecia tubular, with inflated bases, initially polygonal, irregular in transverse section becoming rhombic, bending abruptly in exozones, growing in spiral arrangement from the median axis. Autozoecial diaphragms occurring; hemisepta absent. Autozoecial apertures circular to oval, arranged in regular alternating longitudinal rows. Metazooecia occasionally present. Acanthostyles surrounding autozoecial apertures present, varying in number and size. Autozoecial walls laminated, thick, with dividing hyaline layer in endozone; laminated, merged, containing abundant tubules in exozones (emended after Morozova & Weiss in Morozova et al. 2006).

Remarks.—*Bigeyella* differs from *Lenapora* Ernst & Königshof, 2010, in having acanthostyles and metazooecia.

Stratigraphic and geographic range.—Lower–Upper Devonian (Emsian–Frasnian), Europe, Kazakhstan, Canada.

Bigeyella indigena (Morozova & Weiss in Morozova et al., 2002)

Fig. 4A–C; Table 4.

2002 *Primorella indigena* sp. nov.; Morozova & Weiss in Morozova et al. 2002: 314–315, fig. 6H–J.

2024 *Bigeyella indigena* (Morozova & Weiss in Morozova et al., 2002); Ernst et al. 2024: 15, fig. 3e, f.

Material.—Ten colonies, eighteen thin sections (GIUS 4-292 Gó/Bry A-1a, A-1b, A-1c, A-2a, A-2b, A-2c, A-3a, A-3b, A-4a, A-4b, A-5a, A-6a, A-6b, A-7a, A-8a, A-8b, A-9a, A-10a) from Laskowa Góra Beds, Middle Devonian (upper Givetian) (?*Polygnathus ansatus*–*Schmidtognathus her-*

manni, and *Klapperina disparilis* conodont zones); Górnó, Józefka Hill (road cut, set A), Poland.

Description.—Branched colonies, 0.92–1.90 mm in diameter. Endozone 0.38–0.70 mm wide, exozone 0.24–0.60 mm wide. Axial ratio 0.32–0.50. Autozoecia tubular, budding from distinct branch axis, growing in spiral arrangement from the median axis and bending abruptly in exozones. Diaphragms not observed. Acanthostyles small, with distinct hyaline cores and laminated sheaths. Autozoecial apertures oval, arranged in regular alternating longitudinal rows. Autozoecial walls laminated, 0.02–0.04 mm thick in endozone; laminated, containing tubules in exozone.

Remarks.—*Bigeyella indigena* (Morozova & Weiss in Morozova et al., 2002) differs from *Bigeyella sparsa* Morozova & Weiss in Morozova et al., 2006, in having larger autozoecial apertures (aperture width 0.07–0.13 mm vs. 0.07–0.08 mm in *B. sparsa*).

Stratigraphic and geographic range.—Middle Devonian, upper Givetian of Poland (?*Polygnathus ansatus*–*Schmidtognathus hermanni*, *Klapperina disparilis* conodont zones), Górnó, Józefka Hill (road cut, set A); Upper Devonian, middle Frasnian of Poland: Laskowa Góra Beds. Upper Devonian, ?lower–middle Frasnian, Noravank section, Central Armenia.

Suborder Ptilodictyina Astrova & Morozova, 1956

Family Intraporidae Simpson, 1897

Genus *Intrapora* Hall, 1883

Type species: Intrapora puteolata Hall, 1883; Middle Jeffersonville Limestone, Middle Devonian; Eastern USA.

Emended diagnosis.—Bifoliate colonies consisting of dichotomous branches, leaf-like, frondose. Mesotheca straight or slightly undulating. Autozoecia short, abruptly bending in exozones, Superior hemisepta present or absent. Diaphragms

Table 4. Summary of descriptive statistics for *Bigeyella indigena* (Morozova & Weiss in Morozova et al., 2002) (ten colonies measured). Abbreviations: CV, coefficient of variation; MAX, maximal value; MIN, minimal value; N, number of measurements; SD, sample standard deviation; X, mean.

	N	X	MIN	MAX	SD	CV
Branch width (mm)	7	1.19	0.92	1.90	0.355	29.92
Exozone width (mm)	7	0.35	0.24	0.60	0.135	38.21
Endozone width (mm)	7	0.48	0.38	0.70	0.108	22.53
Axial ratio	7	0.41	0.32	0.50	0.069	16.82
Aperture width (mm)	40	0.10	0.07	0.13	0.019	19.13
Aperture spacing along branch (mm)	20	0.40	0.30	0.48	0.057	14.15
Aperture spacing diagonally (mm)	20	0.25	0.20	0.30	0.025	9.85
Acanthostyle diameter (mm)	20	0.029	0.020	0.040	0.005	17.84

Fig. 5. Fenestrate bryozoans from the Devonian of Poland. **A.** *Laxifenestella* sp., tangential section (GIUS 4-2239 Sk/Bry-XIV-5K-1), from the Dobruchna Member in the bottom part of the Skaly Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone) of Skaly (A₁–A₃). **B, C.** *Juanopora elegans* gen. et sp. nov. from the Wietrznia Beds, upper Givetian (?*Schmidtognathus hermanni* – *Mesotaxis falsiovalis* conodont zones) of Wietrznia-I quarry at Kielce. **B.** Colony fragment, paratype (GIUS 4-260b Wt/Bry-2). **C.** Tangential section showing autozoecial chambers and apertures, holotype (GIUS 4-260b Wt/Br-1a) (C₁–C₅).




Table 5. Summary of descriptive statistics for *Intrapora leunissenii* Ernst, 2008 (four colonies measured). Abbreviations: CV, coefficient of variation; MAX, maximal value; MIN, minimal value; N, number of measurements; SD, sample standard deviation; X, mean.

	N	X	MIN	MAX	SD	CV
Aperture width (mm)	30	0.11	0.09	0.15	0.015	13.58
Aperture spacing along branch (mm)	13	0.49	0.42	0.57	0.048	9.86
Aperture spacing diagonally (mm)	13	0.31	0.27	0.35	0.024	7.53
Metazooecia width (mm)	30	0.05	0.03	0.10	0.016	29.51
Acanthostyle diameter (mm)	30	0.032	0.020	0.043	0.006	18.47

occasionally occurring. Metazooecia abundant, often separating autozoecia. Acanthostyles present or absent.

Remarks.—*Intrapora* Hall, 1883, differs from *Ensiphragma* Astrova in Astrova & Yaroshinskaya, 1968, in arrangement of metazooecia. Metazooecia in *Intrapora* are arranged more or less irregularly, whereas the metazooecia of *Ensiphragma* are arranged in pairs between apertures.

Stratigraphic and geographic range.—Lower Devonian–Mississippian (Emsian–Tournaisian), Europe, North America, and Asia.

Intrapora leunissenii Ernst, 2008

Fig. 4D, E; Table 5.

2008 *Intrapora leunissenii* sp. nov.; Ernst 2008: 357, figs. 25H, 26A–F.

Material.—Four colonies, six thin sections (GIUS 4-2239 Sk/Bry-XIV-5C-1, -5C-2, -5C-3, -5F-1, -5I-1, -5M-1) from Dobruchna Member in the bottom part of the Skały Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone); Skały (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Description.—Bifoliate, branched colonies. Branch cross section lens-shaped. Branches 2.68–2.95 mm wide and 1.63–1.70 mm thick. Mesotheca 0.010–0.015 mm thick, three-layered, with central dark layer and two outer bright layers. Autozoecia long, growing from a mesotheca, semi-circular to trapezoid at the base in cross section, becoming oval to petaloid in the exozone, arranged in 9–15 rows on branches. Autozoecial diaphragms rare, hemisepta absent. Metazooecia small, abundant, separating autozoecia in 1–3 rows, 8–11 surrounding each autozoecial aperture, polygonal to narrowly oval in cross section, often sealed by calcitic skeleton at colony surface. Metazooecial diaphragms abundant, thick. Acanthostyles abundant, 2–5 constantly surrounding each autozoecial aperture and occurring randomly between autozoecia, having narrow hyaline cores and wide laminated sheaths.

Remarks.—*Intrapora leunissenii* Ernst, 2008, is similar to *Intrapora texera* Troitzkaya, 1975, from the Mississippian (Tournaisian) of central Kazakhstan. However, *I. texera* has abundant autozoecial diaphragms and wider branches (branch width about 5 mm vs. 2.68–2.95 mm in *I. leunissenii*).

Stratigraphic and geographic range.—Ahabach-Formation, Middle Devonian, lowermost lower Givetian, Üxheim-Ahütte, Müllertchen quarry, Hillersheim syncline, Eifel,

western Rhenish Massif, Germany. Dobruchna Member in the bottom part of the Skały Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone), Skały (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Order Fenestellida Elias & Condra, 1957

Family Fenestellidae King, 1849

Genus *Rectifenestella* Morozova, 1974

Type species: *Fenestella medvedkensis* Schulga-Nesterenko, 1951; Pennsylvanian (Kasimovian); Russian Platform.

Emended diagnosis.—Reticulate colonies consisting of fine to intermediately robust branches and straight dissepiments. Autozoecia triangular to pentagonal in mid tangential section. Superior hemisepta present; inferior hemisepta absent. Low keel carrying one row of intermediate nodes (modified after Morozova 2001: 45).

Remarks.—*Rectifenestella* Morozova, 1974, differs from *Laxifenestella* Morozova, 1974, in having a pentagonal shape of autozoecia and in the absence of inferior hemisepta, from *Minilya* Crockford, 1944, in having a single row of nodes on the keel instead of two alternating rows in *Minilya*.

Stratigraphic and geographic range.—Devonian–Permian; worldwide.

Rectifenestella sp.

Fig. 4F, G; Table 6.

Material.—Two colonies, four thin sections (GIUS 4-256 GŁ/Bry-1a, -1b, GIUS 4-259 GC/Bry-1a, -1b) from Kadzielnia Limestone Member of the Kowala Formation; Lower Frasnian; *Palmatolepis transitans* Conodont Zone at Bolechowice, Łgawa Hill (set J) and ?*Palmatolepis transitans* Conodont Zone at Kielce, Cmentarna Hill (set B₁), Poland.

Description.—*Exterior:* Reticulate colonies with straight branches, bifurcating, joined by moderately wide dissepiments. Autozoecia arranged in two alternating rows on branches. Autozoecial apertures circular, surrounded by 8 apertural nodes (stellate structure); 3–5 apertures spaced per fenestrule length. Apertural nodes 0.012–0.020 mm in diameter. Fenestrules oval to rectangular, moderately long and narrow. Median keel low. Keel nodes small, intermediately spaced, granular core stellate in shape. Reverse side longitudinally striated.

Interior: Autozoecia triangular to pentagonal in mid-

Table 6. Summary of descriptive statistics for *Rectifenestella* sp. (two colonies measured). Abbreviations: N, number of measurements; X, mean; MIN, minimal value; MAX, maximal value; SD, sample standard deviation; CV, coefficient of variation.

	N	X	MIN	MAX	SD	CV
Branch width (mm)	32	0.28	0.22	0.38	0.037	13.28
Dissepiment width (mm)	30	0.23	0.15	0.36	0.058	24.99
Fenestrule width (mm)	33	0.31	0.22	0.42	0.044	14.15
Fenestrule length (mm)	36	0.74	0.45	1.00	0.104	14.02
Distance between branch centres (mm)	36	0.60	0.47	0.75	0.064	10.62
Distance between dissepiment centres (mm)	39	0.98	0.73	1.20	0.099	10.10
Aperture width (mm)	32	0.08	0.07	0.09	0.008	10.34
Aperture spacing along branch (mm)	32	0.26	0.22	0.32	0.029	10.95
Aperture spacing diagonally (mm)	32	0.23	0.18	0.27	0.022	9.57
Apertures per fenestrule length	40	3.6	3.0	5.0	0.552	15.56
Maximal chamber width (mm)	40	0.13	0.10	0.15	0.010	8.10

tangential section; with well-developed long vestibule; axial wall zigzag; aperture positioned at distal end of chamber. Hemisepta absent. Internal granular skeleton continuous with obverse keel, nodes, peristome and across dissepiments. External laminated skeleton well developed, traversed by abundant microstyles. Microstyles regularly arranged in longitudinal rows on colony reverse surface, 0.02–0.03 mm in diameter. Heterozooecia not observed.

Remarks.—*Rectifenestella* sp. is similar to *Rectifenestella localis* Morozova & Weiss in Morozova et al., 2006, from the Middle Devonian (Eifelian) of Poland, but differs from it in wider branches (branch width 0.22–0.38 mm vs. 0.20–0.25 mm in *R. localis*).

Genus *Laxifenestella* Morozova, 1974

Type species: *Fenestella sarytshevae* Schulga-Nesterenko, 1951, subsequently designated by Morozova (1974); Mississippian (Serpukhovian), Moscow Syncline, Russia.

Emended diagnosis.—Reticulate colonies consisting of relatively wide and thick branches and moderately wide dissepiments. Autozooecia arranged in two rows on the branches. Autozooecial chambers rectangular to pentagonal in mid-tangential section. Axial wall between autozooecial rows weakly undulating. Both superior and inferior hemisepta present. Narrow keel with single row of nodes developed (modified after Morozova 2001: 44).

Remarks.—*Laxifenestella* Morozova, 1974, differs from *Fenestella* Lonsdale, 1839, in the rectangular to pentagonal shape of autozooecia in mid-tangential section and the presence of well-developed hemisepta.

Stratigraphic and geographic range.—Lower Devonian–upper Permian; worldwide.

Laxifenestella sp.

Fig. 5A.

Material.—Single specimen (GIUS 4-2239 Sk/Bry-XIV-5K-1) from Dobruchna Member in the bottom part of the Skąły Formation, uppermost Eifelian (*Polygnathus ensensis*

Conodont Zone); Skąły (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Description.—*Exterior:* Reticulate colonies formed by straight branches joined by relatively wide dissepiments. Branches 0.34–0.37 mm in width, dissepiments 0.28–0.34 mm in width. Fenestrules oval to rectangular, 0.21–0.36 mm in width and 0.47–0.58 mm in length. Autozooecia arranged in two rows on branches. Autozooecial apertures circular, with low peristome; 2 to 3 apertures spaced per fenestrule length. Keel high, nodes inconspicuous. Reverse side not observed.

Interior: Autozooecia relatively long, rectangular in mid-tangential section; with short to moderately long vestibule in longitudinal section. Axial wall between autozooecial rows straight; aperture positioned at distal end of chamber. Superior hemisepta long, proximally curved; inferior hemisepta short or indistinct. External laminated skeleton well-developed on both obverse and reverse side. Heterozooecia not observed.

Remarks.—*Laxifenestella* sp. is superficially similar to *Laxifenestella vera* (Ulrich, 1890) from the Middle Devonian (Givetian) of USA. However, the latter species possesses closely spaced nodes on the keel. This character could not be observed on the present specimen. Material identified as *Laxifenestella vera* has been also described from the Middle Devonian of Mongolia and Altai (Nekhoroshev 1926, 1948; Krasnopeeva 1935).

Genus *Juanopora* nov.

ZooBank LSID: urn:lsid:zoobank.org:act:BB472E06-D35D-447C-9C9E-9189F08745F7

Type species: *Juanopora elegans* gen. et sp. nov., by monotypy; see below.

Etymology: Named in honour of Juan Luis Suárez Andrés (SONIN-GEO, Maliaño, Spain) for his contribution to the knowledge of Palaeozoic bryozoans.

Remarks.—The new genus is mainly characterized by the shape of autozooecia which are long and narrow with rectangular outlines in the middle tangential section. These are unlike those developed in previously documented fenestellid genera. Similar autozooecia are known in the acantho-




Fig. 6. Fenestrate bryozoan *Hemitrypella nodulosa* Ernst et al., 2025, from the Dobruchna Member in the bottom part of the Skały Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone) of Skały, Poland. **A.** Tangential section showing autozoecial chambers and protective superstructure (GIUS 4-2239 Sk/Bry-XIV-5B-1) (A₁–A₃). **B.** Tangential section showing openings of the superstructure (GIUS 4-2239 Sk/Bry-XIV-5P). **C.** Tangential section showing autozoecial apertures and chambers (GIUS 4-2239 Sk/Bry-XIV-5Q) (C₁, C₂).

Table 7. Summary of descriptive statistics for *Juanopora elegans* gen. et sp. nov. (single colony measured). Abbreviations: CV, coefficient of variation; MAX, maximal value; MIN, minimal value; N, number of measurements; SD, sample standard deviation; X, mean.

	N	X	MIN	MAX	SD	CV
Branch width (mm)	5	0.22	0.19	0.25	0.022	9.94
Dissepiment width (mm)	6	0.12	0.09	0.14	0.020	16.67
Fenestrule width (mm)	5	0.30	0.24	0.35	0.040	13.33
Fenestrule length (mm)	3	1.66	1.45	1.88	0.215	12.96
Distance between branch centres (mm)	5	0.49	0.40	0.60	0.078	15.78
Distance between dissepiment centres (mm)	3	1.70	1.60	1.85	0.132	7.78
Aperture width (mm)	5	0.10	0.09	0.11	0.009	9.32
Aperture spacing along branch (mm)	5	0.28	0.23	0.32	0.033	11.60

cladiid genus *Diploporaria* Nickles & Bassler, 1900 (cf. Ernst et al. 2015: 192, fig. 26F–H; *Diploporaria tenella* Wyse Jackson, 1988). However, this genus possesses hemisepta and a pinnate growth form.

Stratigraphic and geographic range.—As for the type species.

Juanopora elegans gen. et sp. nov.

Fig. 5B, C; Table 7.

ZooBank LSID: urn:lsid:zoobank.org:act:B37F7FB6-E7F1-4EDA-AF-CC-5FEB581AE822.

Etymology: From Latin *elegans*, fine; in reference to its elegant appearance and thin morphological elements.

Type material: Holotype, colony fragment (thin section GIUS 4-260b Wt/Br-1a). Paratype, colony fragment (thin section GIUS 4-260b Wt/Bry-2) from the type locality and horizon.

Type locality: Wietrznia-I quarry at Kielce, section Wi-IA, Poland.

Type horizon: Wietrznia Beds (basal set A), upper Givetian (?*Schmidtognathus hermanni*–*Mesotaxis falsiovalis* conodont zones)

Diagnosis.—Reticulate colonies consisting of relatively narrow branches and narrow dissepiments. Fenestrules long and narrow. Autozooezia arranged in two rows on the branches, weakly alternating. Autozooezial chambers rectangular in mid-tangential section, long and narrow. Axial wall between autozooezial rows straight. Hemisepta absent. Keel low; nodes absent.

Description.—*Exterior:* Reticulate colonies formed by straight narrow branches joined by narrow dissepiments. Fenestrules long and narrow, rectangular. Autozooezia arranged in two rows on branches, weakly alternating, often arranged side-by-side (Fig. 5C₃, C₄). Autozooezial apertures circular, with low peristome; 4–6 apertures spaced per fenestrule length. Keel low, nodes absent. Reverse side not observed.

Interior: Autozooezia relatively long, rectangular in mid-tangential section; with short to moderately long vestibule in longitudinal section. Axial wall between autozooezial rows straight; aperture positioned at distal end of chamber. Hemisepta absent. External laminated skeleton well-devel-

oped on both obverse and reverse side. Heterozooezia not observed.

Stratigraphic and geographic range.—Middle Devonian (upper Givetian), Holy Cross Mountains, Poland.

Genus *Hemitrypella* Nekhoroshev, 1948

[= *Neohemitrypa* Schastlivceva in Morozova, 2001]

Type species: *Hemitrypella tubulosa* Nekhoroshev, 1948, by original designation; Middle Devonian; Altai, Russia.

Diagnosis.—Reticulate colonies, conical and conical-tubular shaped, frontal surface exterior. Branches intermediate in width, linear to moderately sinuous, closely or intermediately spaced, dichotomously divided. Two rows of autozooezia per branch, increasing to three rows proximal of branch bifurcations; low straight to sinuous central keel on obverse side of branch with high nodes, composed of core of granular skeleton and sheath of laminar skeleton. Laminar wall extensions of keel nodes fused together forming a fine meshwork of rounded to angular openings, each opening centred over a zooeial aperture in the branch below. Axial wall between autozooezial rows zigzag in tangential sections; autozooezia not strongly inflated laterally, commonly triangular to trapezoid in tangential section; maximum diameter of zooeicia corresponds with either length or height; transverse walls at intermediate or high angle to reverse wall; superior hemisepta absent or weakly developed, other interior structures absent. Small- to large-diameter distal tube typically short, opening frontally or slightly inclined laterally and perhaps distally; apertural peristome present or absent. Heterozooezia not observed. Autozooezial walls of granular material that may be absent on obverse side near apertures; laminar extrazooeial skeleton traversed by small to moderate microstyles (after Ernst et al. 2025).

Remarks.—*Hemitrypella* Nekhoroshev, 1948, differs from *Hemitrypa* Phillips, 1841, in its conical and conical-tubular colony shape and in the triangular to trapezoid shape of autozooezia in tangential section, which are quadrangular or pentagonal in *Hemitrypa*.

Stratigraphic and geographic range.—Lower Devonian

Table 8. Summary of descriptive statistics for *Hemitrypella nodulosa* Ernst et al., 2025 (three colonies measured). Abbreviations: CV, coefficient of variation; MAX, maximal value; MIN, minimal value; N, number of measurements; SD, sample standard deviation; X, mean.

	N	X	MIN	MAX	SD	CV
Branch width (mm)	7	0.23	0.19	0.30	0.042	18.45
Dissepiment width (mm)	8	0.11	0.08	0.16	0.025	22.01
Fenestrule width (mm)	7	0.17	0.14	0.20	0.022	13.15
Fenestrule length (mm)	4	0.40	0.37	0.47	0.047	11.72
Distance between branch centres (mm)	12	0.43	0.35	0.52	0.058	13.74
Distance between dissepiment centres (mm)	4	0.58	0.49	0.63	0.064	11.08
Aperture width (mm)	7	0.07	0.07	0.08	0.005	7.20
Aperture spacing along branch (mm)	8	0.23	0.22	0.26	0.015	6.44
Maximal chamber width (mm)	12	0.12	0.10	0.13	0.010	8.70
Superstructure opening diameter (mm)	13	0.11	0.09	0.13	0.015	13.19
Distance between node centres (mm)	10	0.32	0.26	0.35	0.023	7.22

(Lochkovian)–Mississippian (Tournaisian); Europe, Russia, Kazakhstan, Mongolia, and China.

Hemitrypella nodulosa Ernst et al., 2025

Fig. 6A–C; Table 8.

2025 *Hemitrypella nodulosa* Ernst et al. 2025: 501–503, fig. 6a–h.

Material.—Three colonies, three thin sections (GIUS 4-2239 Sk/Bry-XIV-5P, -5Q, -5B-1) from the Dobruchna Member in the bottom part of the Skąły Formation, Middle Devonian, uppermost Eifelian, *Polygnathus ensensis* Conodont Zone), Skąły, Poland.

Description.—**Exterior:** Reticulate colonies with straight branches joined by dissepiments. Autozoecia arranged in two alternating rows on branches, having circular apertures with low peristomes, 1–2 spaced per length of a fenestrule. Peristomes smooth. Fenestrules oval to slightly rectangular. Openings in the superstructure irregularly shaped, rounded to petaloid, corresponding to positions of apertures, 0.09–0.13 mm in diameter. Reverse colony surface not observed in the present material.

Interior: Autozoecia triangular to trapezoidal in mid-tangential section; low and elongated, with short vestibule in longitudinal section. Axial wall between autozoecial rows zigzag in tangential sections; aperture positioned at distal end of chamber. Hemisepta absent. Internal granular skeleton continuous with obverse keel, nodes, peristome and across dissepiments. External laminated skeleton well developed. Heterozoecia not observed.

Remarks.—*Hemitrypella nodulosa* Ernst et al., 2025, differs from *Hemitrypella tubulosa* Nekhoroshev, 1948, in the wider spacing of branches and dissepiments (average distance between branch centres 0.43 mm vs. 0.33 mm in *H. tubulosa*; average distance between dissepiment centres 0.58 mm vs. 0.39 mm in *H. tubulosa*), and in wider spacing of autozoecial apertures (average distance between centres of adjacent apertures 0.23 mm vs. 0.19 mm in *H. tubulosa*).

Hemitrypella nodulosa Ernst et al., 2025, differs from *Hemitrypella nekhroshevi* Ariunchimeg in Ariunchimeg and Morozova, 1992, from the Middle Devonian (Eifelian) of Mongolia in having shorter fenestrules (fenestrule length 0.37–0.47 mm vs. 0.47–0.60 mm in *H. nekhroshevi*), and in having smaller autozoecial apertures (aperture width 0.07–0.08 mm vs. 0.14 mm in *H. nekhroshevi*).

Stratigraphic and geographic range.—Upper Nims Member of the Junkerberg Formation, Middle Devonian (Eifelian), Brühlborn near Rommersheim, Prüm Syncline in western Rhenish Massif, Germany. Lower Baarley Member of the middle Loogh Formation, Middle Devonian (lower Givetian), “Mühlenwäldchen”, Gerolstein Syncline, Eifel, western Rhenish Massif, Germany. Dobruchna Member in the bottom part of the Skąły Formation, uppermost Eifelian (*Polygnathus ensensis* Conodont Zone), Skąły (trench A dug in 1989; see Woźniak et al. 2022), Poland.

Discussion

As emphasized by Racki et al. (1985) and Morozova et al. (2002), bryozoan distribution in the studied successions was mainly influenced by environmental changes corresponding to deepening pulses (Johnson et al. 1985) and correlative global biotic events (Becker 2025). The partial submergence of the vast *Stringocephalus* Biostromal Bank in the late Givetian during the transgressive cycle IIa was a key ecosystem turnover in the Łysogóry Region and Kostomłoty Zone (Racki et al. 1985, 2022; Fig. 2). As documented by bryozoan-productive sites in the Laskowa Góra Beds (Górno, Laskowa; see Morozova et al. 2002, 2006), the wave of southward expansion of the diverse shelf benthos occurred during the multiphase biotic recovery from the Taghanic global crisis (see also the record of *Canutrypa francqana* in Skompski et al. 2018). This extra-regional immigration is confirmed in this study by the occurrence of *Bigeyella indigena* also in the older Górno fauna (and the occurrence of this species in the Frasnian of Armenia; Ernst et al. 2024), as well as by the new monotypic fenestellid genus described from the Wietrznia site. This finding increases the total number of species in the Wietrznia bryozoan association to seven. Four of these species were new (see Morozova et al. 2006), and, therefore, it can be alternatively assumed that the specific Wietrznia biota is endemic.

The early–middle Frasnian transgressive events were also clearly manifested in the subsequent bryozoan fauna occurring in the fossiliferous Górno succession (7 taxa; Morozova et al. 2002, 2006), associated with very diverse brachiopod fauna (Baliński et al. 2016). Notably, this diversification is not recognized in the coeval shallow-water reef-type habitats, especially in the rich benthic biota of Kadzielnia-type organic buildups. Besides *Rectifenestella*, only tiny problematic lophophorates, the hederellids, are found at the Kadzielnia site.

The present study provides new data concerning the oldest migration to the northern periphery of carbonate platform during the latest Eifelian Kačák (If) event (Fig. 2), recorded in the bryozoan association of the Dobruchna Member. However, taxa lists from both Kiepora (1965, 1973) for the Skąły site and Wyse Jackson in Halamski et al. (2022) for the Miłoszów section suggest that the successive early and middle Givetian (pre-Taghanic) faunas were quite different in composition. This strongly suggests that the pulsed Kačák transgression was associated with a significant, possibly long-term, colonization of the Łysogóry carbonate ramp habitats by this benthic group (Halamski et al. 2022). In particular, the very diverse bryozoan fauna from Miłoszów, comprising at least 20 taxa (Wyse Jackson in Halamski et al. 2022), still awaits a systematic study to provide a basis for estimating the regional effects of the major Taghanic extinction (Ernst 2013).

In the present paper, six species are described from the Dobruchna Member: *Cyclotrypa* sp., *Fistuliphragma gracilis* Ernst, 2008, *Leioclema passitabulatum* Duncan, 1939,

Intrapora leunissenii Ernst, 2008, *Laxifenestella* sp., *Hemitrypella nodulosa* Ernst et al., 2025. From them, *Leioclema passitabulatum*, *Intrapora leunissenii*, and *Hemitrypella nodulosa* are known from the lower Givetian of the Rhenish Massif, whereas *Fistuliphragma gracilis* has originally been described from the Eifelian of the Rhenish Massif. Moreover, *Leioclema passitabulatum* is known from the Lower Devonian of Ukraine (Lochkovian) and Czech Republic (Pragian), as well as from the Middle Devonian (Eifelian) of Michigan, USA. *Fistuliphragma gracilis* was also identified from the Lower to Middle Devonian (Emsian–Eifelian) of Spain (Ernst et al. 2011). Noteworthy, the number of reported bryozoan taxa in the well-known Skafły site is currently extended to 15 (adding 7 encrusting taxa of Ctenostomata after Kiepur 1965: table 1).

In summary, during the IIa and IIc transgressions (Fig. 2), drowning steps of the northern part of the Kielce platform were clearly associated with successive colonization stages of the ramp-type environments. Clearly, the gently inclined slope of the platform, which was subject to dynamic change due to organic growth and synsedimentary tectonic movements, offered several attractive, low-energy niches for lophophorates and echinoderms. This is evident from the thriving populations of diverse, sessile, filter-feeding, benthic organisms, primarily brachiopods and crinoids (Racki et al. 1985; Baliński et al. 2016; Skompski et al. 2018; Halamski et al. 2022). Conversely, a major (in regional perspective; Racki et al. 1985; Racki 1993) latest Givetian IIb deepening episode apparently terminated the development of bryozoan-rich benthic biotas in the Kostomłoty Zone, even though impoverished associations then appeared in the southern Kielce Region (Morozova et al. 2002). The observed replacement of locally abundant and diverse Givetian bank faunas by generally impoverished Frasnian reef-related associations was a remarkable regional trend. It largely corresponds to the worldwide apex of Givetian bryozoans, followed by a major diversity decline in the lineage's evolutionary history (Horowitz et al. 1996; Ernst 2013). However, this turnover probably occurred in two phases in the Holy Cross shelf area (i.e., as a result of the Taghanic and Frasnian crises).

Acknowledgements

AE was supported by the Deutsche Forschungsgemeinschaft (ER 278/4-1 and 2). Catherine Reid (University of Canterbury, Christchurch, UK) and Hans Arne Nakrem (Natural History Museum, Oslo, Norway) are appreciated for their helpful and constructive reviews.

Editor: Andrzej Kaim

References

- Anstey, R.L. and Perry, T.G. 1970. Biometric procedures in taxonomic studies of Paleozoic bryozoans. *Journal of Paleontology* 44: 383–398.
- Ariunchimeg, Y. and Morozova, I.P. 1992. New Paleozoic bryozoans of Mongolia. *Sovmestnaâ Sovetsko-Mongolskaya Paleontologičeskââ Ekspeditsiâ, Trudy* 4: 75–84.
- Astrova, G.G. 1964. New order of paleozoic Bryozoa [in Russian]. *Paleontologičeskij žurnal* 1964 (2): 22–31.
- Astrova, G.G. 1965. Morphology, history of development and system of the Ordovician and Silurian Bryozoa [in Russian]. *Trudy Paleontologičeskogo Instituta Akademii Nauk SSSR* 106: 1–432.
- Astrova, G.G. 1978. The history of development, system, and phylogeny of the Bryozoa: order Trepostomata [in Russian]. *Trudy Paleontologičeskogo Instituta Akademii Nauk SSSR* 169: 1–240.
- Astrova, G.G. and Morozova, I.P. 1956. About systematics of the order Cryptostomata [in Russian]. *Doklady Akademii Nauk SSSR* 110 (4): 661–664.
- Astrova, G.G. and Yaroshinskaya, A.M. 1968. Early Devonian and Eifelian Bryozoa of Salair and Gorno-Altai [in Russian]. *Trudy Tomskogo Gosudarstvennogo Universiteta* 202: 47–62.
- Baliński, A., Racki, G., and Halamski, A. T. 2016. Brachiopods and stratigraphy of the Upper Devonian (Frasnian) succession of the Radlin Syncline (Holy Cross Mountains, Poland). *Acta Geologica Polonica* 66: 125–174.
- Bassler, R.S. 1929. The Permian Bryozoa of Timor. *Paläontologie von Timor* 16: 37–90.
- Bassler, R.S. 1934. Notes on fossil and Recent Bryozoa. *Journal of the Washington Academy of Sciences* 24: 404–408.
- Becker, R.T. 2025. Devonian and lower Carboniferous global events in the central Variscan Orogen. In: U. Linnemann (ed.), *The Variscan Orogen of Central Europe. Regional Geology Reviews*, 889–978. Springer, Cham.
- Boardman, R.S. 1960. Trepostomatous Bryozoa of the Hamilton Group of New York State. *U.S. Geological Survey Professional Papers* 340: 1–87.
- Borg, F. 1926. Studies on Recent cyclostomatous Bryozoa. *Zoologiska Bidrag från Uppsala* 10: 181–507.
- Crockford, J. 1944. Bryozoa from the Permian of Western Australia. Part I. Cyclostomata and Cryptostomata from the north-west basin and Kimberley district. *Proceedings of the New South Wales Linnean Society* 69: 139–175.
- Czaplikowska, I. 1990. *Opracowanie mikropaleontologiczne środkowodevońskich łupków brachiopodowych z Gór Świętokrzyskich*. 79 pp. Unpublished MSc. Thesis, Faculty of Earth Sciences, University of Silesia, Sosnowiec.
- Duncan, H. 1939. Trepostomatous Bryozoa from the Traverse Group of Michigan. *Contributions from the Museum of Paleontology, University of Michigan* 5 (10): 171–270.
- Ehrenberg, C.G. 1831. *Symbolae Physicae, seu Icones et descriptiones Corporum Naturalium novorum aut minus cognitorum, quae ex itineribus per Libyam, Aegyptum, Nubiam, Dongalaam, Syriam, Arabiam et Habessiniam, studia annis 1820–1825, redirent. Pars Zoologica, 4, Animalia Evertebrata exclusis Insectis*. 10 pls. Officina Academica Berolini, Berolini.
- Elias, M.K. and Condra, G.E. 1957. *Fenestella* from the Permian of West Texas. *Geological Society of America Memoir* 70: 1–158.
- Ernst, A. 2008. Non-fenestrate bryozoans from the Middle Devonian of the Eifel (western Rhenish Massif, Germany). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 250: 313–379.
- Ernst, A. 2013. Diversity dynamics and evolutionary patterns of Devonian Bryozoa. *Palaeobiodiversity and Palaeoenvironments* 93: 45–63.
- Ernst, A. and Gorgij, M.N. 2013. Lower Permian bryozoan faunas from Kalmard area, central Iran. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 268: 275–324.
- Ernst, A. and Königshof, P. 2010. Bryozoan fauna and microfacies from a Middle Devonian reef complex (Western Sahara, Morocco). *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 568: 1–91.
- Ernst, A. and May, A. 2009. Bryozoan fauna from the Konęprusy Limestone (Pragian, Lower Devonian) of Zlatý Kůň near Konęprusy (Czech Republic). *Journal of Paleontology* 83 (5): 767–782.
- Ernst, A. and Schröder, S. 2007. Stenolaemate bryozoans from the Middle

- Devonian of the Rhenish Slate Massif (Eifel, Germany). *Neues Jahrbuch für Geologie und Paläontologie* 246: 205–233.
- Ernst, A., Dorsch, T., and Keller, M. 2011. A bryozoan fauna from the Santa Lucia Formation (Lower–Middle Devonian) of Abelgas, Cantabrian Mountains, NW-Spain. *Facies* 57: 301–329.
- Ernst, A., May, A., and Marks, S. 2012. Bryozoans, corals and microfacies of lower Eifelian (Middle Devonian) limestones at Kierspe, Germany. *Facies* 58: 727–758.
- Ernst, A., Wyse Jackson, P. N., and Aretz, M. 2015. Bryozoan fauna from the Mississippian (Viséan) of Roque Redonde (Montagne Noire, southern France). *Geodiversitas* 37: 151–213.
- Ernst, A., Serobyán, V., and Danelian, T. 2024. Biostratigraphic, palaeoenvironmental and palaeobiogeographic implications of bryozoan fauna from the Upper Devonian sequences of Armenia. *Geobios* 85: 10–18.
- Ernst, A., Suárez Andrés, J.L., and Wyse Jackson, P.N. 2025. *Hemistrypa* (Fenestrata, Bryozoa) from the Middle Devonian of the Eifel (western Rhenish Massif) and taxonomic re-evaluation of some allied hemitrypid genera. *Zeitschrift der Deutschen Gesellschaft für Geowissenschaften (Journal of Applied and Regional Geology)* 176: 493–508.
- Gürich, G. 1896. Das Palaeozoicum im Polnischen Mittelgebirge. *Verhandlungen der Russisch-Kaiserlichen Mineralogischen Gesellschaft zu St. Petersburg (series 2)* 32: 1–539.
- Hageman, S.J. 1991a. Approaches to systematic and evolutionary studies of perplexing groups: An example using fenestrata Bryozoa. *Journal of Paleontology* 65: 630–647.
- Hageman, S.J. 1991b. Discrete morphotaxa from a Mississippian fenestrata faunule: presence and implications. In: F.P. Bigey and J.-L. d’Hondt (eds.), *Bryozoaires actuels et fossiles: Bryozoa living and fossil. Bulletin de la Société des Sciences Naturelles de l’Ouest de la France, Mémoire HS 1*: 147–150.
- Halamski, A.T., Baliński, A., Racki, G., Amler, M.R.W., Basse, M., Denayer, J., Dubicka, Z., Filipiak, P., Kondas, M., Krawczyński, W., Mieszkowski, R., Narkiewicz, K., Olempska, E., Wrzolek, T., Wyse Jackson, P.N., Zapalski, M.K., Zatoń, M., and Kozłowski, W. 2022. The pre-Taghanic (Givetian, Middle Devonian) ecosystems of Miłoszów (Holy Cross Mts, Poland). *Annales Societatis Geologorum Poloniae* 92: 323–379.
- Hall, J. 1858. Report on the Geological Survey of Iowa, embracing the results of investigations made during portions of the years 1855, 1856, 1857. *Geological Survey of Iowa, Paleontology 1* (2): 1–724.
- Hall, J. 1883. Bryozoans of the Upper Helderberg and Hamilton groups. *Transactions of the Albany Institute* 10: 145–197.
- Horowitz, A.S., Pachut, J.F., and Anstey, R.L. 1996. Devonian bryozoan diversity, extinctions, and originations. *Journal of Paleontology* 70: 373–380.
- Johnson, J.G., Klapper, G. and Sandberg, C.A. 1985. Devonian eustatic fluctuations in Euramerica. *Geological Society of America Bulletin* 96: 567–587.
- Kiepura, M. 1965. Devonian bryozoans of the Holy Cross Mountains, Poland; Part I, Ctenostomata. *Acta Palaeontologica Polonica* 10: 11–48.
- Kiepura, M. 1973. Devonian bryozoans of the Holy Cross Mountains, Poland. Part 2. Cyclostomata and Cystoporata. *Acta Palaeontologica Polonica* 18: 323–400.
- King, W. 1849. On some families and genera of corals. *Annals and Magazine of the Natural History* 2: 388–390.
- Krasnopeeva, P.S. 1935. Bryozoans of the Middle and Upper Devonian of Altai [in Russian]. *Materialy po geologii Zapadno-Sibirskogo Kraya* 20: 43–84.
- Lonsdale, W. 1839. Corals. In: R.I. Murchison (ed.): *Silurian System*, 675–694. John Murray, London.
- M’Coy, F. 1849. On some new genera and species of Palaeozoic corals and foraminifera. *Annals and Magazine of Natural History* 3: 119–136.
- Ma, J.-Y., Buttler, C.J., and Taylor, P.D. 2014. Cladistic analysis of the “trepotome” suborder Esthonioporina and the systematics of Palaeozoic bryozoans. In: A. Rosso, P.N. Wyse Jackson, and J.S. Porter (eds.), *Bryozoan Studies 2013. Studi Trentini di Scienze Naturali* 94: 153–161.
- Miller, S.A. 1889. *North American Geology and Paleontology*. 664 pp. Western Methodist Book Concern, Cincinnati.
- Morozova, I.P. 1974. Revision of the genus *Fenestella* [in Russian]. *Paleontologičeskij žurnal* 1974 (2): 54–67.
- Morozova, I.P. 2001. Bryozoans of the order Fenestellida [in Russian]. *Trudy Paleontologičeskogo Instituta Rossijskoj Akademii Nauk* 277: 1–176.
- Morozova, I.P., Weis, O.B., and Racki, G. 2002. Emergence and extinction of the Givetian to Frasnian bryozoan faunas in the Kostomłoty facies zone, Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* 47: 307–317.
- Morozova, I.P., Weis, O.B., and Racki, G. 2006. New Devonian and Carboniferous bryozoans of the Holy Cross Mountains (Central Poland). *Paleontologičeskij žurnal* 2006 (5): 58–67.
- Narkiewicz, M., Racki, G., Skompski, S., and Szulczewski, M. 2006. Proceśy i zdarzenia w historii geologicznej Gór Świętokrzyskich. In: S. Skompski and A. Żylińska (eds.), *Materiały konferencyjne, LXXVII Zjazd Naukowy Polskiego Towarzystwa Geologicznego, Ameliówka k. Kielc 28–30 czerwca 2006 r.*, 51–77. Państwowy Instytut Geologiczny, Warszawa.
- Nekhoroshev, V.P. 1926. Middle Devonian bryozoans of northwest Mongolia with a description of the microscopic method for the determination of fenestellids [in Russian]. *Trudy Geologičeskogo muzeja AN SSSR* 1: 1–28.
- Nekhoroshev, V.P. 1948. Devonian bryozoa of the Altai [in Russian]. *Trudy Paleontologičeskogo Instituta Akademii Nauk SSSR* 3: 1–172.
- Nicholson, H.A. 1879. *On the Structure and Affinities of the “Tabulate Corals” of the Paleozoic Period, with Critical Descriptions of Illustrative Species*. 312 pp. William Blackwood and Sons, Edinburgh.
- Nickles, J.M. and Bassler, R.S. 1900. A synopsis of American fossil Bryozoa including bibliography and synonymy. *Bulletin of the United States Geological Survey* 173: 1–663.
- Pajchłowa, M. 1957. The Devonian in the Grzegorzowice-Skały section [in Polish, with English summary]. *Biuletyn Instytutu Geologicznego* 122: 145–254.
- Phillips, J. 1841. *Figures and Descriptions of the Paleozoic Fossils of Cornwall, Devon and West Somerset*. 231 pp. Longman, Brown, Green, Longmans, London.
- Racki, G. 1993. Evolution of the bank to reef complex in the Devonian of the Holy Cross Mountains. *Acta Palaeontologica Polonica* 37 [for 1992]: 87–182.
- Racki, G. 2024. An overlooked contribution to Devonian studies in the Holy Cross Mts: rediscovering Alexei Doronin’s 1893 article on the Kadzielnia Limestone. *Acta Geologica Polonica* 74: e25.
- Racki, G. and Bultynck, P. 1993. Conodont biostratigraphy of the Middle to Upper Devonian boundary beds in the Kielce area of the Holy Cross Mountains. *Acta Geologica Polonica* 43: 1–33.
- Racki, G., Gluchowski, E., and Malec, J. 1985. The Givetian to Frasnian succession at Kostomłoty in the Holy Cross Mts, and its regional significance. *Bulletin of Polish Academy of Sciences. Earth Sciences* 33: 159–171.
- Racki, G., Makowski, I., Mikłás, J., and Gawlik, S. 1993. Brachiopod biofacies in the Frasnian reef-complexes: an example from the Holy Cross Mts, Poland. *Prace Naukowe Uniwersytetu Śląskiego* 1331, *Geologia* 12–13: 64–109.
- Racki, G., Wójcik, K., Halamski, A.T., and Narkiewicz, M. 2022. Middle Devonian Skały Formation in the Holy Cross Mountains (Poland)—formal description and subdivision based on new field data. *Annales Societatis Geologorum Poloniae* 92: 425–444.
- Rominger, C. 1866. Observations on *Chaetetes* and some related genera, in regard to their systematic position; with an appended description of some new species. *Proceedings of the Academy of Natural Sciences of Philadelphia* 18: 113–123.
- Schlüter, C. 1889. Anthozoen des rheinischen Mitteldevon. *Abhandlungen zur geologischen Special-Karte von Preussen und den thüringischen Staaten* 8 (4): 259–465.
- Schulga-Nesterenko, M.I. 1951. Carboniferous Fenestellida of the Russian Platform [in Russian]. *Trudy Paleontologičeskogo Instituta* 32: 1–157.
- Simpson, G.B. 1897. A handbook of the genera of the North American Paleozoic Bryozoa: with an introduction upon the structure of living

- species. In: *14th Annual Report of the State Geologist (of New York) for the Year 1894*, 407–608. James B. Lyon, Albany.
- Skompski, S., Baliński, A., Szulczewski, M., and Zawadzka, I. 2018. Middle/Upper Devonian brachiopod shell concentrations from the intra-shelf basinal carbonates of the Holy Cross Mountains (central Poland). *Acta Geologica Polonica* 68: 607–633.
- Snyder, E.M. 1991. Revised taxonomic procedures and paleoecological applications for some North American Mississippian Fenestellidae and Polyporidae (Bryozoa). *Palaeontographica Americana* 57: 1–275.
- Szulczewski, M. 1995. Depositional evolution of the Holy Cross Mts. (Poland) in the Devonian and Carboniferous—a review. *Geological Quarterly* 3: 471–488.
- Troizkaya, T.D. 1975. Main features of development of bryozoans at the boundary between the Devonian and the Carboniferous in Central Kazakhstan [in Russian]. *Paleontologičeskii žurnal* 1975 (3): 54–69.
- Ulrich, E.O. 1882. American Palaeozoic bryozoa. *Journal of the Cincinnati Society of Natural History* 5: 121–175.
- Ulrich, E.O. 1890. Palaeozoic bryozoa: III. *Geological Survey* 8: 283–688.
- Ulrich, E.O. 1896. Bryozoa. In: C. Eastman (ed.), *Zittel's Textbook of Palaeontology*, Vol. 1, 257–291. Macmillan, London.
- Ulrich, E.O. and Bassler, R.S. 1904. A revision of Palaeozoic Bryozoa. Part II: On genera and species of Trepostomata. *Smithsonian miscellaneous collections* 47: 15–55.
- Vine, G.R. 1884. Fourth report of the Committee consisting of Dr. H.R. Sorby and Mr. G.R. Vine, appointed for the purpose of reporting on fossil Polyzoa. In: *Reports of the 53rd Meeting of the British Association for the Advancement in Sciences*, 161–209. John Murray, London.
- Woźniak, P. 1992. *Środkowodewońskie zespoły ramienionogowe z warstw skalskich w Górach Świętokrzyskich*. 38 pp. Unpublished M.Sc. Thesis, Faculty of Earth Sciences, University of Silesia, Sosnowiec.
- Woźniak, P., Halamski, A.T., and Racki, G. 2022. Cyclic ecological replacement of brachiopod assemblages in the top-Eifelian Dobruchna Brachiopod Shale Member (Skaly Formation) of the Holy Cross Mountains (Poland). *Annales Societatis Geologorum Poloniae* 92: 445–463.
- Wyse Jackson, P.N. 1988. New fenestrate Bryozoa from the Carboniferous of County Fermanagh. *Irish Journal of Earth Sciences* 9: 197–208.
- Yang, J. and Lu, L. 1983. Upper Carboniferous and Lower Permian bryozoa from Kalpin of western Xinjiang. *Paleontologia Cathayana* 1: 259–317.