

## First evidence of attached juveniles in the solutan echinoderm *Pahvanticystis* from the middle Cambrian Weeks Formation (Utah, USA)

Harry J. Savage and Imran A. Rahman

*Acta Palaeontologica Polonica* 70 (1), 2025: 175-178 doi:10.4202/app.01228.2024

The fossil record of the Palaeozoic echinoderm class Soluta suggests they originated in the Miaolingian (middle Cambrian) of Laurentia as permanently attached suspension feeders, demonstrating a stepwise shift towards vagility in successive strata. Here, we report a new specimen of *Pahvanticystis* cf. *utahensis* associated with three putative juveniles interpreted as belonging to the same species. We interpret this as evidence of facultative attachment in juveniles of *Pahvanticystis*, which had not previously been reported in this taxon, but is known in the earlier genus *Castericystis*. Our findings indicate that attachment as a juvenile was more widespread in solutans than previously thought.

Harry J. Savage [[savage\\_harryjames@hotmail.com](mailto:savage_harryjames@hotmail.com); ORCID: <https://orcid.org/0000-0003-3610-5114>], Oxford University Museum of Natural History, Oxford OX1 3PW, UK.

Imran A. Rahman [[imran.rahman@nhm.ac.uk](mailto:imran.rahman@nhm.ac.uk); ORCID: <https://orcid.org/0000-0001-6598-6534>], The Natural History Museum, London SW7 5BD, UK; Oxford University Museum of Natural History, Oxford OX1 3PW, UK.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.