

A new saurodontid fish from the Late Cretaceous of the Western Desert, Egypt

Gebely A. Abu El-Kheir, Ahmed Marzouk, Luigi Capasso, Essam Zahran, Atef M. Kasem, Asmaa Kamel, Mohamed K. Mousa, and Abdelhamid Elbsebeshi

Acta Palaeontologica Polonica 71 (1), 2026: 85-94 doi:10.4202/app.01277.2025

A new genus and species of the extinct family Saurodontidae (Ichthyodectiformes) is described from the Upper Cretaceous deposits of Dakhla Formation, Western Desert, Egypt. The specimen is identified as *Wadiichthys anbaawyi* gen. et sp. nov. The holotype comprises complete skull with maxillae, premaxillae, dentaries and prementary. The specimen was embedded in a compacted mudstone layer of the lower part of Dakhla Formation, Western Desert, Egypt. The new taxon *Wadiichthys anbaawyi* differs from the others saurodontid genera by (i) the dorsal margin of the prementary is longer than the ventral margin. (ii) the longitudinal foramina below each alveolus of the dentary, premaxilla and maxilla; (iii) a small diastema between the teeth; (iv) a convex contact between the rostrodermethmoid and the frontal; (v) the posterior margin of the maxilla is triangular shaped; (vi) the teeth are more deeply embedded in the premaxilla, maxilla and dentary. *Wadiichthys anbaawyi* represents the first formal description of a saurodontid from Africa that contributes to the increase of the knowledge of the diversity and the paleogeographic distribution of this family. The new taxon sheds light on the taxonomy, morphology and evolutionary history of the Cretaceous marine vertebrates of the Tethys Sea. To preserve and digitally document the fossil, a terrestrial laser scanner Trimble TX6 was employed. Each of the five fossil parts was scanned separately and processed with Trimble RealWorks software. This created a high-resolution 3D point cloud in true color, grayscale, and intensity mode. This method protects the fossil from possible damage or loss. It also allows for remote collaboration and morphometric analysis without physical handling.

Key words: Actinopterygii, Ichthyodectiformes, Saurodontidae, Dakhla Formation, Cretaceous, Western Desert, Egypt.

Gebely A. Abu El-Kheir [gebely2006@sci.nvu.edu.eg, ORCID: <https://orcid.org/0000-0001-5905-8790>] and Mohamed K. Mousa [Mohamedkamel.sci@sci.nvu.edu.eg, ORCID: <https://orcid.org/0000-0002-9951-1905>], New Valley Vertebrate Palaeontology Centre, New Valley University, 10th kilometer, Kharga-Assuit road, Kharga, New Valley, Egypt. Ahmed Marzouk [ahmed.marzouk@sci.dmu.edu.eg, ORCID: <https://orcid.org/0009-0006-1111-5163>]

], Essam Zahran [ezahran@sci.dmu.edu.eg], Atef M. Kasem [Kasematef@sci.dmu.edu.eg], and Asmaa Kamel [asmaa.kamel@sci.dmu.edu.eg], Earth Science Department, Faculty of Science, Damanhur University, El Gumhuriya street, Damanhur, Egypt. Luigi Capasso [l.capasso@unich.it, ORCID: <https://orcid.org/0000-0001-6045-3827>], University Museum, 'G. d'Annunzio' University of Chieti–Pescara, Via dei Vestini 31, 66100 Chieti (CH), Italy. Abdelhamid Elbsebeshi [abdelhamid_mohammed@nriag.sci.eg, ORCID: <https://orcid.org/0009-0004-3616-1516>], National Research Institute of Astronomy and Geophysics, Elmarsad street, Helwan, Cairo, Egypt.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,389.2 kB\)](#)