

The first fossil of the extant leatherback sea turtle *Dermochelys coriacea*

Yi-Lu Liaw and Cheng-Hsiu Tsai

Acta Palaeontologica Polonica 71 (2), 2026: 267-271 doi:10.4202/app.01340.2026

The leatherback sea turtle (*Dermochelys coriacea*) is the largest living sea turtle, but their evolutionary history remains poorly known due to the lack of any confirmed fossil record of the extant species. Here we describe a fossil humerus from the Pleistocene of Taiwan. Its preserved morphology shows a close match to the extant leatherback sea turtle, including the humeral head tilted to the intertubercular side, a slightly-curved secondary deltopectoral crest, and the absence of a deep intertubercular fossa. This Pleistocene humerus from Taiwan then presents the first fossil record of the extant leatherback sea turtle worldwide. The record suggests that the modern global migration pattern of the leatherback sea turtle has already included the western North Pacific. Our discovery of the first fossil leatherback sea turtle should encourage more fossil finds to reveal the detailed evolutionary history of the leatherback sea turtle and the ecosystem they utilize.

Yi-Lu Liaw [liawlulu@gmail.com; ORCID: <https://orcid.org/0009-0009-4363-7267>], Department of Life Science, National Taiwan University, Taipei, 10617, Taiwan. Cheng-Hsiu Tsai [whaletsai@ntu.edu.tw; craniata@gmail.com; ORCID: <https://orcid.org/0000-0003-3617-366X>] (corresponding author), Department of Life Science and Institute of Ecology and Evolutionary Biology, National Taiwan University, Taipei, 10617, Taiwan.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(466.8 kB\)](#) |

 [Supplementary file \(433.3 kB\)](#)

