

Note on some *Cranias* from central Poland

Alfred Rosenkrantz

Acta Palaeontologica Polonica 09 (4), 1964: 513-531

Two kinds of *Cranias* have been collected between 193 m. and 207 m. in a boring at Boryszew, central Poland. Most valves represent a new member of the "*Crania tuberculata* group". For members of this group, comprising species from the Maastrichtian, Danian, Paleocene and ?Eocene, a new subgenus of *Crania* - *C. (Danocrania)* n.subg. is erected with *Crania tuberculata* Nilsson as type species. The Polish representatives are named: *C. (Danocrania) polonica* n.sp. and *C. (Danocrania) polonica semicostulata* n.subsp. The subspecies shows a peculiar outer sculpture very similar to that shown by a *C. (Danocrania)* from the upper Middle Danian of Faxe (Denmark). The remaining *Cranias* all belong to the subgenus *C. (Isocrania)*, known only in association with *C. (Danocrania)* in Danian strata. The Polish form is related to *C. (Isocrania) posselti* Rkz. from the Upper Danian of Denmark. The age of the *Crania*-bearing strata at Boryszew is estimated to be upper Middle Danian. If the strata are younger, their Danian fossil content, e.g. the *Cranias*, *Tylocidaris masoviensis* Kongiel and *Stereocidaris rosenkrantzi* (Ravn), must be considered as derived fossils. A ventral valve of the Upper Senonian *C. (Isocrania) paucicostata* Bosquet in Lower Paleocene strata at Bochońnica shows that derived fossils may occur in the Lower Paleocene of central Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.