

Revision of the flexible crinoid genus *Ammonocrinus* and a new hypothesis on its life mode

Jan Bohatý


Acta Palaeontologica Polonica 56 (3), 2011: 615-639 doi: <http://dx.doi.org/10.4202/app.2010.0020>

The lecanocrinid *Ammonocrinus* (Flexibilia) is newly interpreted based on new material from the Middle Devonian of the Rhenish Massif (Eifel and Bergisches Land, Germany). The species have echinoid-like tubercles on the attachment and on the column, which bear articulated spines. The intraspecific variability of the column is discussed for three facies-controlled morphotypes, herein classified as standard “exposed–” or “encased roller-type” and the rare “settler-type”. New specimens have floating transitions between different plate sculpturing and between those individuals with none or one to several columnals with herein termed “lateral columnal enclosure extensions” on the proximal–most, barrel-like dististele and the following mesistele, which is solely distinguished by these extensions. Based on this interpretation, *Ammonocrinus kongieli* is evaluated as a subjective junior synonym of *Ammonocrinus sulcatus*. The latter species was first recognised from the Eifel (Germany). “*Ammonocrinus wachtbergensis*”, from the upper Eifelian of the Eifel, is declared a subjective junior synonym of *Ammonocrinus doliiformis*. The first nearly complete specimen of *Ammonocrinus kerdreoletensis* is described from the lower Eifelian of Vireux–Molhain (southern Ardennes, France). Two new species are described: *Ammonocrinus jankei* sp. nov. and *Ammonocrinus leunissenii* sp. nov. A functional morphologic trend in perfecting the crown encasement by continuous modification of the lateral columnal enclosure extensions of the mesistele from the Eifelian to the Givetian, indicates a vagile benthic “predator”–driven evolution of ammonicrinids in the Eifel area. Several ammonicrinid species are herein defined as spined soft–bottom dwellers, feeding in low–intensity current water, possibly through a self–produced water flow. The first known postmortem encrusting epizoans on ammonicrinid endoskeletons are reported.

Key words: Crinoidea, Flexibilia, *Ammonocrinus*, Devonian, Eifel, Rhenish Massif, Germany.

Jan Bohatý [jan.bohaty@mail.com], Institut für Geologie und Mineralogie der Universität zu Köln, Zülpicher Str. 49a, D-50674 Köln, Germany.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,917.8 kB\)](#)