

## ***Xenoxylon* synecology and palaeoclimatic implications for the Mesozoic of Eurasia**

Changhwan Oh, Marc Philippe, and Kyungsik Kim


*Acta Palaeontologica Polonica* 60 (1), 2015: 245-256 doi: <http://dx.doi.org/10.4202/app.2012.0132>

The distribution of fossil wood genera has been demonstrated to be an effective proxy for Mesozoic terrestrial climates. In this study, we investigated the phytocoenoses, which were associated with *Xenoxylon* confirmed to be a marker for a cool and/or wet climate in a boreal hemisphere (i.e., *Xenoxylon*-phytocoenoses) during the Mesozoic, using specimens of fossil wood. It was confirmed that *Xenoxylon* co-occurs more often with some wood genera than with others. For example, *Protocedroxylon*, a wood that is most likely related to the Pinaceae, is the genus most often associated with *Xenoxylon*-phytocoenoses. Although *Taxodioxygen* is also found in *Xenoxylon*-phytocoenoses, it is not found, however, as consistently as *Protocedroxylon*. The distribution and diversity of *Xenoxylon*-phytocoenoses changed throughout the Mesozoic. During the Late Triassic and Late Cretaceous, *Xenoxylon*-phytocoenoses had low diversity and were restricted to higher palaeolatitudes during the Late Cretaceous. However, during the Early to Middle Jurassic, *Xenoxylon*-phytocoenoses were distributed much farther south, while their diversity concomitantly increased sharply. From the Late Jurassic to the Early Cretaceous, the distribution of *Xenoxylon*-phytocoenoses moved northward in Europe and even more so in East Asia. The changes in the distribution of *Xenoxylon*-phytocoenoses are in agreement with changes in both global and regional climates. Our results also demonstrated that, within the *Xenoxylon* distribution range, the corresponding phytocoenoses were differentiated along a latitudinal gradient and according to the global climate change patterns during the Mesozoic.

**Key words:** Fossil wood, *Xenoxylon*-phytocoenoses, palaeoecology, palaeoclimate, Mesozoic, Eurasia.

Changhwan Oh [[bluegaia@kopri.re.kr](mailto:bluegaia@kopri.re.kr)] Division of Polar Earth-System Sciences, Korea Polar Research Institute, 26 Songdomirae-ro, Yeonsu-gu, Incheon, 406-840, Republic of Korea; Marc Philippe [[marc.philippe@univ-lyon1.fr](mailto:marc.philippe@univ-lyon1.fr)], Paléobotanique, Université Claude Bernard-Lyon 1 and UMR5276 of the CNRS, F69622, France; Kyungsik Kim [[kksik@jbnu.ac.kr](mailto:kksik@jbnu.ac.kr)], Department of Life Sciences, Chonbuk National University, 570-752, Republic of Korea.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(527.5 kB\)](#) |

 [Supplementary file \(199.2 kB\)](#)