

***Catopsalis* (Multituberculata) from Asia and North America and the problem of taeniolabidid dispersal in the Late Cretaceous**

Zofia Kielan-Jaworowska and Robert E. Sloan

Acta Palaeontologica Polonica 24 (2), 1979: 187-197

Djadochtatherium Simpson, 1925 is a junior synonym of *Catopsalis* Cope, 1882. Known species of *Catopsalis* (*Djadochtatherium* included) form a morphological sequence, beginning with Asian ?Late Santonian and/or Early Campanian *C. matthewi*, and ?Middle Campanian *C. catopsaloides*, through the following North American species: Late Maastrichtian *C. joyneri*, Paleocene *C. foliatus*, *C. utahensis*, *C. fissidens* and *C. cagartensis*. Morphological changes involved an increase in size, a reduction of the number of upper premolars, relative reduction of the size of the lower fourth premolars, and an increase of the number of cusps on the molars. It is believed that *Catopsalis*, and possibly the family Taeniolabididae originated in Asia, developed there during the ?Late Santonian-?Middle Campanian and spread to North America during the Late Campanian or Early Maastrichtian.

Key words: Cretaceous, migrations, Multituberculata, Paleocene, Taeniolabididae.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.