

## Early astogeny and relationships of the Ordovician rhabdomesine bryozoans

Jerzy Dzik

*Acta Palaeontologica Polonica* 37 (1), 1992: 37-54

A diverse Ordovician assemblage of juvenile bryozoan colonies, with external phosphatic coatings allowing chemical extraction from the rock, shows a morphologic series probably corresponding to evolutionary step-by-step suppression in the formation of the external colony wall. Its formation was initially delayed leading to formation of the common bud (advanced corynotrypids). Then the permanent double-walled colony organization developed along the colony margin (*Flabellotrypa*-like). In the following evolutionary stage, the external wall was secreted only at the conical creeping stage immediately following ancestrula (typical rhabdomesines). With subsequent evolution the external wall completely disappeared, and the double-walled ancestrula grew vertically (advanced rhabdomesines and phylloporinids). This kind of the early astogeny characterizes also the oldest phyllodictyid cryptostome *Prophyllodictya*, studied in serial peels. Flabellotrypidae fam. n., *Mojczatrypa halysitoides* gen. et sp. n., Kielceporidae fam. n., *Kielcepora ornata* gen. et sp. n., *Ojlepora* gen. n., and *Kielanopora gracilis* gen. et sp. n. are proposed.

**Key words:** astogeny, evolution, Ordovician, Bryozoa

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.