

## Anatomical distinctions of the Mesozoic lingulide brachiopods

Gertruda Biernat and Christian C. Emig

*Acta Palaeontologica Polonica* 38 (1-2), 1993: 1-20

The long held view that *Lingula* represents an extremely bradytelic lineage is questioned. Examination of Mesozoic lingulides has shown that they significantly differ from their Recent relatives *Lingula* and *Glottidia* in having longer lophophoral cavities, shorter ventral canals, better developed posterior adductor muscles, and less acute umbones. Morphological characters of the shell interior are needed to identify members of the Lingulidae, not solely external shell characteristics. The apparent evolutionary tendency towards a reduction of the volume of the lophophoral cavity contradicts the traditional view that the 'living fossil' *Lingula* has survived without significant morphological change since the Paleozoic. Actually the today living lingulide genera probably arose in the early Cenozoic. A new inarticulate brachiopod genus, *Lingularia* is introduced, with three new species, Middle Triassic *L. siberica*, Middle Jurassic *L. sirnilis*, and Cretaceous *L. smirnome*.

**Key words:** lingulides, Brachiopoda, shell morphology, taxonomy, evolution, Paleozoic-Recent.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.