

New Early Jurassic sawflies from Luxembourg : the oldest record of Tenthredinoidea (Hymenoptera: 'Symphyta')

André Nel, Julián F. Petrulevicius, and Michel Henrotay

Acta Palaeontologica Polonica 49 (2), 2004: 283-288

Pseudoxyelocerus bascharagensis gen. et sp. nov., the oldest representative of the Tenthredinoidea and Xyelotomidae, based on a single forewing, and an enigmatic 'Symphyta' family *incertae sedis*, based on a hindwing, are described from the Toarcian of Luxembourg. The relationships of the genera currently included in Xyelotomidae are briefly discussed. The genera *Undatoma*, *Liaotoma*, *Leridatoma*, and *Davidsmithia* have the unique apomorphy in the wing venation of the Tenthredinoidea minus Xyelotomidae. The Xyelotomidae is probably a paraphyletic family. Only a phylogenetic analysis will help to solve these problems. There is no evidence supporting the previous assignments of the fossil genera *Vitimilarva* and *Kuengilarva* to the family Xyelotomidae.

Key words: Hymenoptera, Tenthredinoidea, Xyelotomidae, *Pseudoxyelocerus*, Toarcian, Luxembourg.

André Nel [anel@mnhn.fr] and Julián F. Petrulevicius (CONICET) [levicius@mnhn.fr], Laboratoire d'Entomologie (CNRS UMR 5146), Muséum National d'Histoire Naturelle, 45 rue Buffon, F-75005 Paris, France; M. Henrotay, 8 rue de Bouny, B-4624, Romsée, Belgique.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.