

Late Eocene scyliorhinid sharks from the Trans-Urals, Russia

Tatiana Malyshkina

Acta Palaeontologica Polonica 51 (3), 2006: 465-475

Priabonian deposits from two localities, Kurgan and Derney, in the Trans-Urals (Western Siberia) have yielded numerous selachian teeth. The carcharhiniform family Scyliorhinidae is represented by three species, including two new: *Premontreia uralica* sp. nov., *Fountizia zhelezkoi* sp. nov., and *F. pattersoni* (Cappetta, 1976). Both genera are recorded from the West-Siberian Basin for the first time. Presumably all three species (including the one morphologically mimicking *F. pattersoni*, a species recorded from Ypresian of England) are endemic for the Trans-Urals region (contrary to the cosmopolitan lamniforms recorded previously from the area. The known distribution pattern of extant scyliorhinids supports the probability of endemism of the cat sharks from the West-Siberian Basin, which has also been isolated geographically from the Peri-Tethys during the Late Eocene marine regression. The peculiar local environmental conditions due to the Priabonian climatic cooling in the Boreal realm might have also facilitated the speciation.

Key words: : Elasmobranchii, Carcharhiniformes, Scyliorhinidae, *Premontreia*, *Fountizia*, Eocene, West-Siberian Basin.

Tatiana Malyshkina malyshkina@igg.uran.ru, Institute of Geology and Geochemistry of the Russian Academy of Sciences Urals Branch, Pochtovyi per. 7, Ekaterinburg, 620151, Russia

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.