

A review of the endocerid cephalopod *Protocyptendoceras* from the Floian (Lower Ordovician) of the Eastern Cordillera, Argentina.

Marcela Cichowolski

Acta Palaeontologica Polonica 54 (1), 2009: 99-109 doi:<http://dx.doi.org/10.4202/app.2009.0110>

Ordovician cephalopods from Western Gondwana are still poorly known and published data is in need of systematic and stratigraphical revision. In this work, the endocerid proterocameroceratid *Protocyptendoceras* from the Floian of the Eastern Cordillera of Argentina is revised using new material from La Ciénaga, 5 km upstream from Purmamarca. The species *P. corvalani* and *P. teichert* are considered to be junior synonyms of *P. fuenzalidae*, which is redescribed in detail and reillustrated. Specimens preserve adapical parts of isolated siphuncles, related here to the presence of endocones in a posterior portion. Evidently *P. fuenzalidae* had a nekto-benthic mode of life in a shallow water environment, oriented horizontally. Its palaeobiogeographical affinities are mainly with Floian proterocameroceratids from Eastern Gondwana, such as *Anthoceras decorum* and the related genera *Lobendoceras*, *Ventroloboceras*, and *Notocycloceras*.

Key words: Cephalopoda, Endocerida, Proterocameroceratidae, Floian, Ordovician, Eastern Cordillera, Argentina.

Marcela Cichowolski [mcicho@gl.fcen.uba.ar], Área de Paleontología, Departamento de Ciencias Geológicas, Universidad de Buenos Aires, Ciudad Universitaria, Pab. II, Intendente Güiraldes 2160 C1428EGA, Buenos Aires, Argentina. CONICET.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

