

## Morphology and ontogeny of some Middle Ordovician gonambonitid brachiopods from Baltoscandia

Anna A. Madison

*Acta Palaeontologica Polonica* 63 (3), 2018: 585-594 doi:<https://doi.org/10.4202/app.00488.2018>

The Volkhovian–Kunda boundary deposits of the Leningrad Region contain two closely related genera, *Antigonambonites* and *Anchigonites*. The latter genus was previously known only by valve moulds; here its shell exterior and interior, and ontogeny are described in detail based on the collection of well-preserved disarticulated valves of *Anchigonites conulus*. *Antigonambonites* and *Anchigonites* share similar developmental and morphological features, and ecological strategies including the attachment by cementation by a “pedicle tube” and are thus possibly phylogenetically related. A new diagnosis for the genus *Anchigonites* and description of *A. conulus* are provided; other species of *Anchigonites* are briefly revised. The moulds of outer epithelial cells are described for the first time for the class Strophomenata.

**Key words:** Brachiopoda, Gonambonitidae, morphology, ontogeny, Ordovician, Russia.

Anna A. Madison [[sunnyannmad@yahoo.com](mailto:sunnyannmad@yahoo.com)], Borissiak Paleontological Institute of Russian Academy of Sciences, 123, Profsoyuznaya ul., Moscow, 117997 Russia.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.