

## Sequence of post-moult exoskeleton hardening preserved in a trilobite mass moult assemblage from the Lower Ordovician Fezouata Konservat-Lagerstätte, Morocco

Harriet B. Drage, Thijs R.A. Vandenbroucke, Peter Van Roy, and Allison C. Daley

*Acta Palaeontologica Polonica* 64 (2), 2019: 261-273 doi:<https://doi.org/10.4202/app.00582.2018>

Euarthropods have a tough exoskeleton that provides crucial protection from predation and parasitism. However, this is restrictive to growth and must be periodically moulted. The moulting sequence is well-known from extant arthropods, consisting of: (i) the long inter-moult stage, in which no changes occur to the hardened exoskeleton; (ii) the pre-moult stage where the old exoskeleton is detached and the new one secreted; (iii) exuviation, when the old exoskeleton is moulted; and (iv) the post-moult stage during which the new exoskeleton starts as soft, thin, and partially compressed and gradually hardens to the robust exoskeleton of the inter-moult stage. Trilobite fossils typically consist of inter-moult carcasses or moulted exuviae, but specimens preserving the post-moult stage are rare. Here we describe nine specimens assigned to *Symphysurus ebbestadi* representing the first group of contemporaneous fossils collected that preserve all key stages of the moulting process in one taxon, including the post-moult stage. They were collected from a single lens in the Tremadocian part of the Fezouata Shale Formation, Morocco. Based on cephalic displacement and comparison to other trilobite moults, one specimen appears to represent a moulted exoskeleton. Four specimens are typical inter-moult carcasses. Four others are wrinkled and flattened, with thin exoskeletons compared to inter-moult specimens, and are considered post-moult individuals. These *S. ebbestadi* specimens illuminate the preservation and morphology of the post-moulting stage, characterised by strong anterior-posterior exoskeleton wrinkling, as well as overall body flattening and reduced visibility of thoracic articulations. Being found in the same lens, these specimens likely represent the first preserved in-the-act mass moulting event. The displayed sequence of moulting suggests the moulting process in trilobites was comparable to modern arthropods, and conserved within euarthropod evolutionary history.

**Key words:** Trilobita, mass moult, soft-shelled, post-moult, moulting, exoskeleton, Ordovician, Morocco.

Harriet B. Drage [[harriet.drage@zoo.ox.ac.uk](mailto:harriet.drage@zoo.ox.ac.uk)], Department of Zoology, University of Oxford, South Parks Road, OX1 3PS, UK; and Oxford University Museum of Natural History, Parks Road, OX1 3PW, UK.

Thijs R.A. Vandenbroucke [[thijs.vandenbroucke@ugent.be](mailto:thijs.vandenbroucke@ugent.be)] and Peter Van Roy [[Peter.VanRoy@ugent.be](mailto:Peter.VanRoy@ugent.be)], Department of Geology, Ghent University, Campus

Sterre, 9000 Ghent, Belgium. Allison C. Daley [[allison.daley@unil.ch](mailto:allison.daley@unil.ch)], Institute of Earth Sciences, University of Lausanne, Géopolis, CH-1015 Lausanne, Switzerland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



[Full text \(788.8 kB\)](#) |



[Supplementary file \(88.6 kB\)](#)