

## New species of mirid insects and their importance for the higher classification of plant bugs

Junggon Kim, Artur Taszakowski, Aleksander Herczek, Marzena Zmarzły and Sunghoon Jung  
*Acta Palaeontologica Polonica* 68 (1) 2023: 75-83 doi:<https://doi.org/10.4202/app.00991.2022>

The detailed morphological study based on the findings of well-preserved fossil specimens in Eocene Baltic amber revealed two new species, *Metoisops michalskii* Kim, Taszakowski, and Herczek sp. nov. and *Metoisops popovi* Kim, Taszakowski, and Jung sp. nov. The morphological information of the extinct genus *Metoisops* including diagnoses, descriptions of new species, and a species key are provided. The divided fourth antennal segment is depicted. The tribal transfer of *Metoisops* from Electromyiommini to Gigantometopini is proposed based on major morphological characters, five-six femoral trichobothria, presence of a deep incision between calli, and the structure of parameres. The need for a phylogenetic revision of the internal classification within Isometopinae is also discussed.

**Key words:** Hemiptera, Heteroptera, Cimicomorpha, Miridae, Isometopinae, classification, jumping tree bugs, Baltic amber.

Junggon Kim [[thesv12@gmail.com](mailto:thesv12@gmail.com); ORCID: <https://orcid.org/0000-0003-0594-7618>], 1719 Gyebaek-ro, Jung-gu, Daejeon, Korea. Artur Taszakowski [[artur.taszakowski@us.edu.pl](mailto:artur.taszakowski@us.edu.pl); ORCID: <https://orcid.org/0000-0002-0885-353X>], Aleksander Herczek [[aleksander.herczek@us.edu.pl](mailto:aleksander.herczek@us.edu.pl); ORCID: <https://orcid.org/0000-0001-6047-5268>], and Marzena Zmarzły [[marzena.zmarzly@us.edu.pl](mailto:marzena.zmarzly@us.edu.pl); ORCID: <https://orcid.org/0000-0002-0631-6079>], Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences, University of Silesia in Katowice, Bankowa 9, 40-007 Katowice, Poland. Sunghoon Jung [[jung@cnu.ac.kr](mailto:jung@cnu.ac.kr); ORCID: <https://orcid.org/0000-0001-6086-0326>] (corresponding author), Laboratory of Systematic Entomology, Department of Applied Biology, College of Agriculture and life Sciences, Chungnam National University, 99, Daehak-ro, Daejeon, Korea and Department of Smart Agriculture Systems, College of Agriculture and Life Sciences, Chungnam National University, 99, Daehak-ro, Daejeon, Korea.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

 [Full text \(1,154.2 kB\)](#)