

## Structure of the K-T mammal radiation in North America: Speculations on turnover rates and trophic structure

J. David Archibald

*Acta Palaeontologica Polonica* 28 (1-2), 1983: 7-17

Preliminary results from Montana and New Mexico (USA) indicate that appearance rates for mammal species increase from the latest Cretaceous (Lancian) into the early Paleocene (Puercan) with a slight decline in to the middle Paleocene (Torrejonian). Disappearance rates decline over the same interval and equal appearance rates by the Torrejonian. These results are the opposite of those found by Van Valen (1978). Some groups (condylarthrans) have more frequent speciation events and shorter species durations than suggested by Stanley (1979). The overall appearance rate for the K-T interval is very similar to that for the Paleocene-Eocene transition (both about  $1 \text{ sp./}10^4 \text{ yrs.}$ ). The overall disappearance rate for the K-T interval ( $1.4 \text{ sp./}10^4 \text{ yrs.}$ ) is markedly higher than for the Paleocene-Eocene transition ( $0.5 \text{ sp./}10^4 \text{ yrs.}$ ). With the extinction of dinosaurs, "guilds" of small mammals are re-established and new "guilds" form after about  $2.5 \times 10^4$  years. Appearance rates are highest during guild establishment.

**Key words:** Mesozoic mammals, Cenozoic mammal radiation, Cretaceous-Tertiary boundary, turnover rates, trophic structure.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](http://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.