

Late Frasnian Athyridida (Brachiopoda) from Poland and the Late Devonian biotic turnover

Tatjana A. Grunt and Grzegorz Racki

Acta Palaeontologica Polonica 43 (2), 1998: 361-378

Late Frasnian representatives of the order Athyridida from the Holy Cross Mountains, Poland, support the idea that the Laurussian basins were the places of origin and radiation of the subfamilies Athyridinae and Meristinae during the middle and early late Paleozoic. At least three new species have been identified from two localities (Łgawa Hill and Kowala) in the Gałęzice Syncline. Of these, one was probably endemic (*Merista rhenanensis* sp. n.; maybe also ?*Zonathyris* sp. A), and two (*Athyris postconcentrica* sp. n. and *Pachyplaxoides postgyralea* gen. et sp. n.) were more widely distributed in this part of the Laurussian shelf, being known also from the East European Platform and Rheinisches Schiefergebirge, respectively. This confirms an intermediate biogeographic position of the Holy Cross Mountains area, belonging to an important centre of brachiopod origin and diversification. In contrast to other articulate brachiopods, athyridids reveal a higher rate of diversification, especially at the species (and partly also generic) level, during the global Kellwasser Crisis.

Key words: Brachiopoda, Athyridida, taxonomy, biostratigraphy, biogeography, phylogeny, mass-extinction, Kellwasser Crisis, Frasnian, Famennian, Devonian, Poland.

Tatjana A. Grunt [lenin33@paleo.msk.su], Paleontological Institute, Russian Academy of Sciences, Profsoyuznaya Str., 123, 117647 Moscow, Russia. Grzegorz Racki [racki@us.edu.pl], Katedra Paleontologii i Stratygrafii, Uniwersytet Śląski, ul. Będzińska 60, PL-41-200 Sosnowiec, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

