

***Agastograptus*, a synonym of *Plectograptus* (Retiolitidae, Graptolithina)**

Anna Kozłowska-Dawidziuk

Acta Palaeontologica Polonica 47 (3), 2002: 459-467

Agastograptus robustus Obut and Zaslavskaya, 1983, the type species of the genus *Agastograptus* is herein recognized as a species of *Plectograptus*. Large proximal lateral orifices are one of the significant characters of *Plectograptus*. *P. robustus* differs from the type species, *Plectograptus macilentus* Törnquist, 1887, in possessing paired apertural processes. Other species of *Agastograptus* have been assigned to three different genera: *Spinograptus*, *S. clathrospinosis* (Eisenack, 1951), *S. munchi* (Eisenack, 1951), *Neogothograptus*, *N. balticus* (Eisenack, 1951), and *Cometograptus*, *C. nevadensis* (Berry and Murphy, 1975). The main diagnostic feature used for *Agastograptus*, the spinoreticular paired apertural processes, is recognized as a species feature, characteristic mostly for *Spinograptus*, whereas the generic features are the arrangement of the proximal end, ventral walls, and ancora sleeve of the rhabdosome. Therefore the genus *Agastograptus* is a synonym of *Plectograptus*.

Key words: Graptolithina, Retiolitidae, Agastograptus, Silurian, taxonomy

Anna Kozłowska-Dawidziuk [akd@twarda.pan.pl], Instytut Paleobiologii PAN, ul. Twarda 51/55, PL-00-818 Warszawa, Poland.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see creativecommons.org), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.