

## The oldest known Miocene snake fauna from Central Europe: Merkur-North locality, Czech Republic

Martin Ivanov

*Acta Palaeontologica Polonica* 47 (3), 2002: 513-534

The Early Miocene Merkur-North locality (MN 3a) represents the oldest known Miocene ophidian locality in Europe east of Germany. The snake assemblage is characterised by high species diversity and includes the following families: Boidae (*Bavarioboa* sp.), Colubridae (*Coluber dolnicensis*, *Coluber suevicus*, *Coluber caspioides*, cf. *Elaphe* sp., *Natrix merkurensis* sp. nov., *Natrix sansaniensis*), Elapidae (Elapidae gen. et sp. indet.), and Viperidae (*Vipera* sp.-"*Vipera aspis*" complex). Fossils of the extinct species, *Coluber dolnicensis*, *Coluber suevicus*, and *Natrix sansaniensis*, represent their earliest known occurrences. The cranial elements of *C. suevicus* and *N. sansaniensis* are described for the first time. Discoveries of cf. *Elaphe* sp. may represent the earliest fossil member of the genus *Elaphe*. Elapidae gen. et sp. indet. probably represents the oldest known member of the cobras.

**Key words:** Snakes, Boidae, Colubridae, Elapidae, Viperidae, Early Miocene, Czech Republic

Martin Ivanov [[mivanov@mzm.cz](mailto:mivanov@mzm.cz)], Department of Geology and Palaeontology, Moravian Museum, Zelný trh 6, 659 37 Brno, Czech Republic.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see [creativecommons.org](https://creativecommons.org)), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.