

## The earliest annelids: Lower Cambrian polychaetes from the Sirius Passet Lagerstätte, Peary Land, North Greenland

Simon Conway Morris and John S. Peel *Acta Palaeontologica Polonica* 53 (1), 2008: 137-148 doi:http://dx.doi.org/10.4202/app.2008.0110

Apart from the Phyllopod Bed of the Burgess Shale (Middle Cambrian) polychaete annelids are practically unknown from any of the Cambrian Lagerstätten. This is surprising both because their diversity in the Burgess Shale is considerable, while to date the Chengjiang Lagerstätte which is equally impressive in terms of faunal diversity has no reliable records of any annelids. Here we describe, on the basis of about 40 specimens, *Phragmochaeta* canicularis gen. et sp. nov. from the Lower Cambrian Sirius Passet Lagerstätte of Peary Land, North Greenland. This makes it by far the oldest known polychaete, with a likely age of lower to middle Atdabanian, The body consists of approximately 20 segments, each bearing notochaetae and neurochaetae. The former appeared to have formed a felt-like covering on the dorsum, whilst the neurochaetae projected obliquely to the longitudinal axis. Apart from minor differences in chaetal size at either end there is no other tagmosis. Details of the head are obscure, and presence of palps, tentacles and eyes are conjectural. Jaws appear to have been absent. The gut was straight, and flanked by massive longitudinal musculature. P. canicularis was evidently benthic, propelling itself on the neurochaetae, with the dorsal neurochaetae conferring protection. Its stratigraphic position and generalized appearance are consistent with P. canicularis being primitive, but the phylogenetic relationships within the polychaetes remain problematic, principally because of paucity of relevant morphological information.

**Key words:** Annelida, Polychaeta, Phagmochaeta, metazoan evolution, Cambrian, Sirius Passet Lagerstätte, Greenland.

Simon Conway Morris <a href="mailto:sc113@esc.cam.ac.uk">sc113@esc.cam.ac.uk</a>, Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EQ, England; John. S. Peel <a href="mailto:john.peel@pal.uu.se">john.peel@pal.uu.se</a>, Department of Earth Sciences and Museum of Evolution, Uppsala University, Villavägen 16, SE-752 36 Uppsala, Sweden.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (for details please see <u>creativecommons.org</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Full text (2,194.4 kB)